

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International BureauExp Mail EV741777646US
USAN 10/010,742

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C12N 15/12, 15/62, C07K 14/47, 16/18, A61K 35/12, 38/17, 39/00, A61P 35/00, G01N 33/53, C12Q 1/68, A61K 48/00	A2	(11) International Publication Number: WO 00/61756 (43) International Publication Date: 19 October 2000 (19.10.00)
(21) International Application Number: PCT/US00/09688 (22) International Filing Date: 10 April 2000 (10.04.00) (30) Priority Data: 09/288,950 9 April 1999 (09.04.99) US 09/346,327 2 July 1999 (02.07.99) US (71) Applicant (for all designated States except US): CORIXA CORPORATION [US/US]; Suite 200, 1124 Columbia Street, Seattle, WA 98104 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): REED, Steven, G. [US/US]; 2843 - 122nd Place NE, Bellevue, WA 98005 (US). XU, Jiangchun [US/US]; 15805 SE 43rd Place, Bellevue, WA 98006 (US). DILLON, Davin, C. [US/US]; 21607 NE 24th Street, Redmond, WA 98053 (US). (74) Agents: POTTER, Jane, E.R.; Seed Intellectual Property Law Group PLLC, Suite 6300, 701 Fifth Avenue, Seattle, WA 98104-7092 (US) et al.		(81) Designated States: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: COMPOUNDS FOR IMMUNOTHERAPY AND DIAGNOSIS OF BREAST CANCER AND METHODS FOR THEIR USE		
(57) Abstract Compounds and methods for the treatment and diagnosis of breast cancer are provided. The inventive compounds include polypeptides containing at least a portion of a breast tumor protein. Vaccines and pharmaceutical compositions for immunotherapy of breast cancer comprising such polypeptides, or polynucleotides encoding such polypeptides, are also provided, together with polynucleotides for preparing the inventive polypeptides.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

COMPOUNDS FOR IMMUNOTHERAPY AND DIAGNOSIS
OF BREAST CANCER AND METHODS FOR THEIR USE

5 TECHNICAL FIELD

The present invention relates generally to compositions and methods for the treatment and diagnosis of breast cancer. The invention is more particularly related to polypeptides comprising at least a portion of a protein that is preferentially expressed in breast tumor tissue and to polynucleotides encoding such polypeptides.

10 Such polypeptides and polynucleotides may be used in vaccines and pharmaceutical compositions for treatment of breast cancer. Additionally such polypeptides and polynucleotides may be used in the immunodiagnosis of breast cancer.

BACKGROUND OF THE INVENTION

15 Breast cancer is a significant health problem for women in the United States and throughout the world. Although advances have been made in detection and treatment of the disease, breast cancer remains the second leading cause of cancer-related deaths in women, affecting more than 180,000 women in the United States each year. For women in North America, the life-time odds of getting breast
20 cancer are now one in eight.

No vaccine or other universally successful method for the prevention or treatment of breast cancer is currently available. Management of the disease currently relies on a combination of early diagnosis (through routine breast screening procedures) and aggressive treatment, which may include one or more of a variety of
25 treatments such as surgery, radiotherapy, chemotherapy and hormone therapy. The course of treatment for a particular breast cancer is often selected based on a variety of prognostic parameters, including an analysis of specific tumor markers. See, e.g., Porter-Jordan and Lippman, *Breast Cancer* 8:73-100 (1994). However, the use of established markers often leads to a result that is difficult to interpret, and the high

mortality observed in breast cancer patients indicates that improvements are needed in the treatment, diagnosis and prevention of the disease.

Accordingly, there is a need in the art for improved methods for therapy and diagnosis of breast cancer. The present invention fulfills these needs and
5 further provides other related advantages.

SUMMARY OF THE INVENTION

The present invention provides compounds and methods for immunotherapy of breast cancer. In one aspect, isolated polypeptides are provided
10 comprising at least an immunogenic portion of a breast tumor protein or a variant of said protein that differs only in conservative substitutions and/or modifications, wherein the breast tumor protein comprises an amino acid sequence encoded by a polynucleotide comprising a sequence selected from the group consisting of (a)
nucleotide sequences recited in SEQ ID NOS: 3, 10, 17, 24, 45-52, 55-67, 72, 73, 89-
15 97, 102 and 107, (b) complements of said nucleotide sequences and (c) sequences that hybridize to a sequence of (a) or (b) under moderately stringent conditions. In specific embodiments, the isolated polypeptides of the present invention comprise an amino acid sequence of SEQ ID NO: 98, 99 or 101.

In related aspects, isolated polynucleotides encoding the above
20 polypeptides are provided. In specific embodiments, such polynucleotides comprise sequences provided in SEQ ID NOS: 3, 10, 17, 24, 45-52 and 55-67, 72, 73, 89-97, 102 and 107. The present invention further provides expression vectors comprising the above polynucleotides and host cells transformed or transfected with such expression vectors. In preferred embodiments, the host cells are selected from the
25 group consisting of *E. coli*, yeast and mammalian cells.

In another aspect, the present invention provides fusion proteins comprising a first and a second inventive polypeptide or, alternatively, an inventive polypeptide and a known breast antigen.

The present invention also provides pharmaceutical compositions
30 comprising at least one of the above polypeptides, or a polynucleotide encoding such a polypeptide, and a physiologically acceptable carrier, together with vaccines

comprising at least one or more such polypeptide or polynucleotide in combination with a non-specific immune response enhancer. Pharmaceutical compositions and vaccines comprising one or more of the above fusion proteins are also provided.

In related aspects, pharmaceutical compositions for the treatment of breast cancer comprising at least one polypeptide and a physiologically acceptable carrier are provided, wherein the polypeptide comprises an immunogenic portion of a breast tumor protein or a variant thereof, the breast tumor protein being encoded by a polynucleotide comprising a sequence selected from the group consisting of: (a) nucleotide sequences recited in SEQ ID NOS: 1, 2, 4-9, 11-16, 18-23, 25-44, 53, 54, 68-71, 74-88 and 103-106, (b) complements of said nucleotide sequences, and (c) sequences that hybridize to a sequence of (a) or (b) under moderately stringent conditions. The invention also provides vaccines for the treatment of breast cancer comprising such polypeptides in combination with a non-specific immune response enhancer, together with pharmaceutical compositions and vaccines comprising at least one polynucleotide comprising a sequence provided in SEQ ID NOS: 1, 2, 4-9, 11-16, 18-23, 25-44, 53, 54, 68-71, 74-88 and 103-106.

In yet another aspect, methods are provided for inhibiting the development of breast cancer in a patient, comprising administering an effective amount of at least one of the above pharmaceutical compositions and/or vaccines.

The present invention also provides methods for immunodiagnosis of breast cancer, together with kits for use in such methods. In one specific aspect of the present invention, methods are provided for detecting breast cancer in a patient, comprising: (a) contacting a biological sample obtained from a patient with a binding agent that is capable of binding to one of the above polypeptides; and (b) detecting in the sample a protein or polypeptide that binds to the binding agent. In preferred embodiments, the binding agent is an antibody, most preferably a monoclonal antibody.

In related aspects, methods are provided for monitoring the progression of breast cancer in a patient, comprising: (a) contacting a biological sample obtained from a patient with a binding agent that is capable of binding to one

of the above polypeptides; (b) determining in the sample an amount of a protein or polypeptide that binds to the binding agent; (c) repeating steps (a) and (b); and comparing the amounts of polypeptide detected in steps (b) and (c).

Within related aspects, the present invention provides antibodies, preferably monoclonal antibodies, that bind to the inventive polypeptides, as well as diagnostic kits comprising such antibodies, and methods of using such antibodies to inhibit the development of breast cancer.

The present invention further provides methods for detecting breast cancer comprising: (a) obtaining a biological sample from a patient; (b) contacting the sample with a first and a second oligonucleotide primer in a polymerase chain reaction, at least one of the oligonucleotide primers being specific for a polynucleotide that encodes one of the above polypeptides; and (c) detecting in the sample a DNA sequence that amplifies in the presence of the first and second oligonucleotide primers. In a preferred embodiment, at least one of the oligonucleotide primers comprises at least about 10 contiguous nucleotides of a polynucleotide comprising a sequence selected from the group consisting of SEQ ID NOS: 1-97, 100 and 102-107.

In a further aspect, the present invention provides a method for detecting breast cancer in a patient comprising: (a) obtaining a biological sample from the patient; (b) contacting the sample with an oligonucleotide probe specific for a polynucleotide that encodes one of the above polypeptides; and (c) detecting in the sample a polynucleotide sequence that hybridizes to the oligonucleotide probe. Preferably, the oligonucleotide probe comprises at least about 15 contiguous nucleotides of a polynucleotide comprising a sequence selected from the group consisting of SEQ ID NOS: 1-97, 100 and 102-107.

In related aspects, diagnostic kits comprising the above oligonucleotide probes or primers are provided.

These and other aspects of the present invention will become apparent upon reference to the following detailed description. All references disclosed herein

are hereby incorporated by reference in their entirety as if each was incorporated individually.

5 BRIEF DESCRIPTION OF THE DRAWINGS AND SEQUENCE IDENTIFIERS

Figs. 1A and B show the specific lytic activity of a first and a second B511S-specific CTL clone, respectively, measured on autologous LCL transduced with B511s (filled squares) or HLA-A3 (open squares).

- 10 SEQ ID NO: 1 is the determined 3'cDNA sequence of 1T-5120
- SEQ ID NO: 2 is the determined 3'cDNA sequence of 1T-5122
- SEQ ID NO: 3 is the determined 3'cDNA sequence of 1T-5123
- SEQ ID NO: 4 is the determined 3'cDNA sequence of 1T-5125
- SEQ ID NO: 5 is the determined 3'cDNA sequence of 1T-5126
- 15 SEQ ID NO: 6 is the determined 3'cDNA sequence of 1T-5127
- SEQ ID NO: 7 is the determined 3'cDNA sequence of 1T-5129
- SEQ ID NO: 8 is the determined 3'cDNA sequence of 1T-5130
- SEQ ID NO: 9 is the determined 3'cDNA sequence of 1T-5133
- SEQ ID NO: 10 is the determined 3'cDNA sequence of 1T-5136
- 20 SEQ ID NO: 11 is the determined 3'cDNA sequence of 1T-5137
- SEQ ID NO: 12 is the determined 3'cDNA sequence of 1T-5139
- SEQ ID NO: 13 is the determined 3'cDNA sequence of 1T-5142
- SEQ ID NO: 14 is the determined 3'cDNA sequence of 1T-5143
- SEQ ID NO: 15 is the determined 5'cDNA sequence of 1T-5120
- 25 SEQ ID NO: 16 is the determined 5'cDNA sequence of 1T-5122
- SEQ ID NO: 17 is the determined 5'cDNA sequence of 1T-5123
- SEQ ID NO: 18 is the determined 5'cDNA sequence of 1T-5125
- SEQ ID NO: 19 is the determined 5'cDNA sequence of 1T-5126
- SEQ ID NO: 20 is the determined 5'cDNA sequence of 1T-5127
- 30 SEQ ID NO: 21 is the determined 5'cDNA sequence of 1T-5129
- SEQ ID NO: 22 is the determined 5'cDNA sequence of 1T-5130

- SEQ ID NO: 23 is the determined 5'cDNA sequence of 1T-5133
SEQ ID NO: 24 is the determined 5'cDNA sequence of 1T-5136
SEQ ID NO: 25 is the determined 5'cDNA sequence of 1T-5137
SEQ ID NO: 26 is the determined 5'cDNA sequence of 1T-5139
5 SEQ ID NO: 27 is the determined 5'cDNA sequence of 1T-5142
SEQ ID NO: 28 is the determined 5'cDNA sequence of 1T-5143
SEQ ID NO: 29 is the determined 5'cDNA sequence of 1D-4315
SEQ ID NO: 30 is the determined 5'cDNA sequence of 1D-4311
SEQ ID NO: 31 is the determined 5'cDNA sequence of 1E-4440
10 SEQ ID NO: 32 is the determined 5'cDNA sequence of 1E-4443
SEQ ID NO: 33 is the determined 5'cDNA sequence of 1D-4321
SEQ ID NO: 34 is the determined 5'cDNA sequence of 1D-4310
SEQ ID NO: 35 is the determined 5'cDNA sequence of 1D-4320
SEQ ID NO: 36 is the determined 5'cDNA sequence of 1E-4448
15 SEQ ID NO: 37 is the determined 5'cDNA sequence of 1S-5105
SEQ ID NO: 38 is the determined 5'cDNA sequence of 1S-5110
SEQ ID NO: 39 is the determined 5'cDNA sequence of 1S-5111
SEQ ID NO: 40 is the determined 5'cDNA sequence of 1S-5116
SEQ ID NO: 41 is the determined 5'cDNA sequence of 1S-5114
20 SEQ ID NO: 42 is the determined 5'cDNA sequence of 1S-5115
SEQ ID NO: 43 is the determined 5'cDNA sequence of 1S-5118
SEQ ID NO: 44 is the determined 5'cDNA sequence of 1T-5134
SEQ ID NO: 45 is the determined 5'cDNA sequence of 1E-4441
SEQ ID NO: 46 is the determined 5'cDNA sequence of 1E-4444
25 SEQ ID NO: 47 is the determined 5'cDNA sequence of 1E-4322
SEQ ID NO: 48 is the determined 5'cDNA sequence of 1S-5103
SEQ ID NO: 49 is the determined 5'cDNA sequence of 1S-5107
SEQ ID NO: 50 is the determined 5'cDNA sequence of 1S-5113
SEQ ID NO: 51 is the determined 5'cDNA sequence of 1S-5117
30 SEQ ID NO: 52 is the determined 5'cDNA sequence of 1S-5112

- SEQ ID NO: 53 is the determined cDNA sequence of 1013E11
SEQ ID NO: 54 is the determined cDNA sequence of 1013H10
SEQ ID NO: 55 is the determined cDNA sequence of 1017C2
SEQ ID NO: 56 is the determined cDNA sequence of 1016F8
5 SEQ ID NO: 57 is the determined cDNA sequence of 1015F5
SEQ ID NO: 58 is the determined cDNA sequence of 1017A11
SEQ ID NO: 59 is the determined cDNA sequence of 1013A11
SEQ ID NO: 60 is the determined cDNA sequence of 1016D8
SEQ ID NO: 61 is the determined cDNA sequence of 1016D12
10 SEQ ID NO: 62 is the determined cDNA sequence of 1015E8
SEQ ID NO: 63 is the determined cDNA sequence of 1015D11
SEQ ID NO: 64 is the determined cDNA sequence of 1012H8
SEQ ID NO: 65 is the determined cDNA sequence of 1013C8
SEQ ID NO: 66 is the determined cDNA sequence of 1014B3
15 SEQ ID NO: 67 is the determined cDNA sequence of 1015B2
SEQ ID NO: 68-71 are the determined cDNA sequences of previously identified
antigens
SEQ ID NO: 72 is the determined cDNA sequence of JJ9434
SEQ ID NO: 73 is the determined cDNA sequence of B535S
20 SEQ ID NO: 74-88 are the determined cDNA sequence of previously identified
antigens
SEQ ID NO: 89 is the determined cDNA sequence of B534S
SEQ ID NO: 90 is the determined cDNA sequence of B538S
SEQ ID NO: 91 is the determined cDNA sequence of B542S
25 SEQ ID NO: 92 is the determined cDNA sequence of B543S
SEQ ID NO: 93 is the determined cDNA sequence of P501S
SEQ ID NO: 94 is the determined cDNA sequence of B541S
SEQ ID NO: 95 is an extended cDNA sequence for 1016F8 (also referred to as
B511S)
30 SEQ ID NO: 96 is an extended cDNA sequence for 1016D12 (also referred to as

B532S)

SEQ ID NO: 97 is an extended cDNA sequence for 1012H8 (also referred to as B533S)

SEQ ID NO: 98 is the predicted amino acid sequence for B511S

5 SEQ ID NO: 99 is the predicted amino acid sequence for B532S

SEQ ID NO: 100 is the determined full-length cDNA sequence for P501S

SEQ ID NO: 101 is the predicted amino acid sequence for P501S

SEQ ID NO: 102 is the determined cDNA sequence of clone #19605, also referred to as 1017C2, showing no significant homology to any known gene

10 SEQ ID NO: 103 is the determined 3' end cDNA sequence for clone #19599, showing homology to the Tumor Expression Enhanced gene

SEQ ID NO: 104 is the determined 5' end cDNA sequence for clone #19599, showing homology to the Tumor Expression Enhanced gene

15 SEQ ID NO: 105 is the determined cDNA sequence for clone #19607, showing homology to Stromelysin-3

SEQ ID NO: 106 is the determined cDNA sequence for clone #19601, showing homology to Collagen

SEQ ID NO: 107 is the determined cDNA sequence of clone #19606, also referred to as B546S, showing no significant homology to any known gene

20

DETAILED DESCRIPTION OF THE INVENTION

As noted above, the present invention is generally directed to compositions and methods for the immunotherapy and diagnosis of breast cancer. The inventive compositions are generally isolated polypeptides that comprise at least
25 a portion of a breast tumor protein. Also included within the present invention are molecules (such as an antibody or fragment thereof) that bind to the inventive polypeptides. Such molecules are referred to herein as "binding agents."

In particular, the subject invention discloses isolated polypeptides comprising at least a portion of a human breast tumor protein, or a variant thereof,
30 wherein the breast tumor protein includes an amino acid sequence encoded by a

polynucleotide molecule including a sequence selected from the group consisting of: nucleotide sequences recited in SEQ ID NOS: 1-97, 100 and 102-107, the complements of said nucleotide sequences, and variants thereof. In certain specific embodiments, the inventive polypeptides comprise an amino acid sequence selected
5 from the group consisting of sequences provided in SEQ ID NO: 98, 99 and 101, and variants thereof. As used herein, the term "polypeptide" encompasses amino acid chains of any length, including full length proteins, wherein the amino acid residues are linked by covalent peptide bonds. Thus, a polypeptide comprising a portion of one of the above breast proteins may consist entirely of the portion, or the portion
10 may be present within a larger polypeptide that contains additional sequences. The additional sequences may be derived from the native protein or may be heterologous, and such sequences may be immunoreactive and/or antigenic.

As used herein, an "immunogenic portion" of a human breast tumor protein is a portion that is capable of eliciting an immune response in a patient
15 inflicted with breast cancer and as such binds to antibodies present within sera from a breast cancer patient. Such immunogenic portions generally comprise at least about 5 amino acid residues, more preferably at least about 10, and most preferably at least about 20 amino acid residues. Immunogenic portions of the proteins described herein may be identified in antibody binding assays. Such assays may generally be
20 performed using any of a variety of means known to those of ordinary skill in the art, as described, for example, in Harlow and Lane, *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory, Cold Spring Harbor, NY, 1988. For example, a polypeptide may be immobilized on a solid support (as described below) and contacted with patient sera to allow binding of antibodies within the sera to the
25 immobilized polypeptide. Unbound sera may then be removed and bound antibodies detected using, for example, ¹²⁵I-labeled Protein A. Alternatively, a polypeptide may be used to generate monoclonal and polyclonal antibodies for use in detection of the polypeptide in blood or other fluids of breast cancer patients. Methods for preparing and identifying immunogenic portions of antigens of known sequence are well known

in the art and include those summarized in Paul, *Fundamental Immunology*, 3rd ed., Raven Press, 1993, pp. 243-247.

The term "polynucleotide(s)," as used herein, means a single or double-stranded polymer of deoxyribonucleotide or ribonucleotide bases and includes DNA and corresponding RNA molecules, including HnRNA and mRNA molecules, both sense and anti-sense strands, and comprehends cDNA, genomic DNA and recombinant DNA, as well as wholly or partially synthesized polynucleotides. An HnRNA molecule contains introns and corresponds to a DNA molecule in a generally one-to-one manner. An mRNA molecule corresponds to an HnRNA and DNA molecule from which the introns have been excised. A polynucleotide may consist of an entire gene, or any portion thereof. Operable anti-sense polynucleotides may comprise a fragment of the corresponding polynucleotide, and the definition of "polynucleotide" therefore includes all such operable anti-sense fragments.

The compositions and methods of the present invention also encompass variants of the above polypeptides and polynucleotides. A polypeptide "variant," as used herein, is a polypeptide that differs from the recited polypeptide only in conservative substitutions and/or modifications, such that the therapeutic, antigenic and/or immunogenic properties of the polypeptide are retained. In a preferred embodiment, variant polypeptides differ from an identified sequence by substitution, deletion or addition of five amino acids or fewer. Such variants may generally be identified by modifying one of the above polypeptide sequences, and evaluating the antigenic properties of the modified polypeptide using, for example, the representative procedures described herein. Polypeptide variants preferably exhibit at least about 70%, more preferably at least about 90% and most preferably at least about 95% identity (determined as described below) to the identified polypeptides.

As used herein, a "conservative substitution" is one in which an amino acid is substituted for another amino acid that has similar properties, such that one skilled in the art of peptide chemistry would expect the secondary structure and hydropathic nature of the polypeptide to be substantially unchanged. In general, the

following groups of amino acids represent conservative changes: (1) ala, pro, gly, glu, asp, gln, asn, ser, thr; (2) cys, ser, tyr, thr; (3) val, ile, leu, met, ala, phe; (4) lys, arg, his; and (5) phe, tyr, trp, his.

5 Variants may also, or alternatively, contain other modifications, including the deletion or addition of amino acids that have minimal influence on the antigenic properties, secondary structure and hydrophobic nature of the polypeptide. For example, a polypeptide may be conjugated to a signal (or leader) sequence at the N-terminal end of the protein which co-translationally or post-translationally directs transfer of the protein. The polypeptide may also be conjugated to a linker or other
10 sequence for ease of synthesis, purification or identification of the polypeptide (e.g., poly-His), or to enhance binding of the polypeptide to a solid support. For example, a polypeptide may be conjugated to an immunoglobulin Fc region.

A nucleotide "variant" is a sequence that differs from the recited nucleotide sequence in having one or more nucleotide deletions, substitutions or
15 additions. Such modifications may be readily introduced using standard mutagenesis techniques, such as oligonucleotide-directed site-specific mutagenesis as taught, for example, by Adelman et al. (*DNA*, 2:183, 1983). Nucleotide variants may be naturally occurring allelic variants, or non-naturally occurring variants. Variant nucleotide sequences preferably exhibit at least about 70%, more preferably at least
20 about 80% and most preferably at least about 90% identity (determined as described below) to the recited sequence.

The antigens provided by the present invention include variants that are encoded by DNA sequences which are substantially homologous to one or more of the DNA sequences specifically recited herein. "Substantial homology," as used
25 herein, refers to DNA sequences that are capable of hybridizing under moderately stringent conditions. Suitable moderately stringent conditions include prewashing in a solution of 5X SSC, 0.5% SDS, 1.0 mM EDTA (pH 8.0); hybridizing at 50°C-65°C, 5X SSC, overnight or, in the event of cross-species homology, at 45°C with 0.5X SSC; followed by washing twice at 65°C for 20 minutes with each of 2X, 0.5X and
30 0.2X SSC containing 0.1% SDS. Such hybridizing DNA sequences are also within

the scope of this invention, as are nucleotide sequences that, due to code degeneracy, encode an immunogenic polypeptide that is encoded by a hybridizing DNA sequence.

Two nucleotide or polypeptide sequences are said to be "identical" if the sequence of nucleotides or amino acid residues in the two sequences is the same when aligned for maximum correspondence as described below. Comparisons between two sequences are typically performed by comparing the sequences over a comparison window to identify and compare local regions of sequence similarity. A "comparison window" as used herein, refers to a segment of at least about 20 contiguous positions, usually 30 to about 75, more preferably 40 to about 50, in which a sequence may be compared to a reference sequence of the same number of contiguous positions after the two sequences are optimally aligned.

Optimal alignment of sequences for comparison may be conducted using the Megalign program in the Lasergene suite of bioinformatics software (DNASTAR, Inc., Madison, WI), using default parameters. This program embodies several alignment schemes described in the following references: Dayhoff, M.O. (1978) A model of evolutionary change in proteins – Matrices for detecting distant relationships. In Dayhoff, M.O. (ed.) *Atlas of Protein Sequence and Structure*, National Biomedical Research Foundation, Washington DC Vol. 5, Suppl. 3, pp. 345-358; Hein J. (1990) *Unified Approach to Alignment and Phylogenies* pp. 626-645 *Methods in Enzymology* vol. 183, Academic Press, Inc., San Diego, CA; Higgins, D.G. and Sharp, P.M. (1989) Fast and sensitive multiple sequence alignments on a microcomputer *CABIOS* 5:151-153; Myers, E.W. and Muller W. (1988) Optimal alignments in linear space *CABIOS* 4:11-17; Robinson, E.D. (1971) *Comb. Theor* 11:105; Santou, N. Nes, M. (1987) The neighbor joining method. A new method for reconstructing phylogenetic trees *Mol. Biol. Evol.* 4:406-425; Sneath, P.H.A. and Sokal, R.R. (1973) *Numerical Taxonomy – the Principles and Practice of Numerical Taxonomy*, Freeman Press, San Francisco, CA; Wilbur, W.J. and Lipman, D.J. (1983) Rapid similarity searches of nucleic acid and protein data banks *Proc. Natl. Acad. Sci. USA* 80:726-730.

Preferably, the “percentage of sequence identity” is determined by comparing two optimally aligned sequences over a window of comparison of at least 20 positions, wherein the portion of the polynucleotide sequence in the comparison window may comprise additions or deletions (i.e. gaps) of 20 percent or less, usually 5 to 15 percent, or 10 to 12 percent, as compared to the reference sequences (which does not comprise additions or deletions) for optimal alignment of the two sequences. The percentage is calculated by determining the number of positions at which the identical nucleic acid bases or amino acid residue occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the reference sequence (i.e. the window size) and multiplying the results by 100 to yield the percentage of sequence identity.

Also included in the scope of the present invention are alleles of the genes encoding the nucleotide sequences recited herein. As used herein, an “allele” or “allelic sequence” is an alternative form of the gene which may result from at least one mutation in the nucleic acid sequence. Alleles may result in altered mRNAs or polypeptides whose structure or function may or may not be altered. Any given gene may have none, one, or many allelic forms. Common mutational changes which give rise to alleles are generally ascribed to natural deletions, additions, or substitutions of nucleotides. Each of these types of changes may occur alone or in combination with the others, one or more times in a given sequence.

For breast tumor polypeptides with immunoreactive properties, variants may alternatively be identified by modifying the amino acid sequence of one of the above polypeptides, and evaluating the immunoreactivity of the modified polypeptide. For breast tumor polypeptides useful for the generation of diagnostic binding agents, a variant may be identified by evaluating a modified polypeptide for the ability to generate antibodies that detect the presence or absence of breast cancer. Such modified sequences may be prepared and tested using, for example, the representative procedures described herein.

The breast tumor proteins of the present invention, and polynucleotide molecules encoding such proteins, may be isolated from breast tumor tissue using any

of a variety of methods well known in the art. Polynucleotide sequences corresponding to a gene (or a portion thereof) encoding one of the inventive breast tumor proteins may be isolated from a breast tumor cDNA library using a subtraction technique as described in detail below. Examples of such DNA sequences are provided in SEQ ID NOS: 1- 97, 100 and 102-107. Partial polynucleotide sequences thus obtained may be used to design oligonucleotide primers for the amplification of full-length polynucleotide sequences in a polymerase chain reaction (PCR), using techniques well known in the art (see, for example, Mullis et al., *Cold Spring Harbor Symp. Quant. Biol.*, 51:263, 1987; Erlich ed., *PCR Technology*, Stockton Press, NY, 1989). Once a polynucleotide sequence encoding a polypeptide is obtained, any of the above modifications may be readily introduced using standard mutagenesis techniques, such as oligonucleotide-directed site-specific mutagenesis as taught, for example, by Adelman et al. (*DNA*, 2:183, 1983).

The breast tumor polypeptides disclosed herein may also be generated by synthetic or recombinant means. Synthetic polypeptides having fewer than about 100 amino acids, and generally fewer than about 50 amino acids, may be generated using techniques well known to those of ordinary skill in the art. For example, such polypeptides may be synthesized using any of the commercially available solid-phase techniques, such as the Merrifield solid-phase synthesis method, where amino acids are sequentially added to a growing amino acid chain (see, for example, Merrifield, *J. Am. Chem. Soc.* 85:2149-2146, 1963). Equipment for automated synthesis of polypeptides is commercially available from suppliers such as Perkin Elmer/Applied BioSystems Division (Foster City, CA), and may be operated according to the manufacturer's instructions.

Alternatively, any of the above polypeptides may be produced recombinantly by inserting a polynucleotide sequence that encodes the polypeptide into an expression vector and expressing the protein in an appropriate host. Any of a variety of expression vectors known to those of ordinary skill in the art may be employed to express recombinant polypeptides of this invention. Expression may be achieved in any appropriate host cell that has been transformed or transfected with an

expression vector containing a polynucleotide molecule that encodes a recombinant polypeptide. Suitable host cells include prokaryotes, yeast and higher eukaryotic cells. Preferably, the host cells employed are *E. coli*, yeast or a mammalian cell line, such as CHO cells. The polynucleotide sequences expressed in this manner may
5 encode naturally occurring polypeptides, portions of naturally occurring polypeptides, or other variants thereof.

In general, regardless of the method of preparation, the polypeptides disclosed herein are prepared in an isolated, substantially pure form (*i.e.*, the polypeptides are homogenous as determined by amino acid composition and primary
10 sequence analysis). Preferably, the polypeptides are at least about 90% pure, more preferably at least about 95% pure and most preferably at least about 99% pure. In certain preferred embodiments, described in more detail below, the substantially pure polypeptides are incorporated into pharmaceutical compositions or vaccines for use in one or more of the methods disclosed herein.

15 In a related aspect, the present invention provides fusion proteins comprising a first and a second inventive polypeptide or, alternatively, a polypeptide of the present invention and a known breast tumor antigen, together with variants of such fusion proteins.

A polynucleotide sequence encoding a fusion protein of the present
20 invention is constructed using known recombinant DNA techniques to assemble separate polynucleotide sequences encoding the first and second polypeptides into an appropriate expression vector. The 3' end of a polynucleotide sequence encoding the first polypeptide is ligated, with or without a peptide linker, to the 5' end of a polynucleotide sequence encoding the second polypeptide so that the reading frames
25 of the sequences are in phase to permit mRNA translation of the two DNA sequences into a single fusion protein that retains the biological activity of both the first and the second polypeptides.

A peptide linker sequence may be employed to separate the first and the second polypeptides by a distance sufficient to ensure that each polypeptide folds
30 into its secondary and tertiary structures. Such a peptide linker sequence is

incorporated into the fusion protein using standard techniques well known in the art. Suitable peptide linker sequences may be chosen based on the following factors: (1) their ability to adopt a flexible extended conformation; (2) their inability to adopt a secondary structure that could interact with functional epitopes on the first and second polypeptides; and (3) the lack of hydrophobic or charged residues that might react with the polypeptide functional epitopes. Preferred peptide linker sequences contain Gly, Asn and Ser residues. Other near neutral amino acids, such as Thr and Ala may also be used in the linker sequence. Amino acid sequences which may be usefully employed as linkers include those disclosed in Maratea et al., *Gene* 40:39-46, 1985; Murphy et al., *Proc. Natl. Acad. Sci. USA* 83:8258-8262, 1986; U.S. Patent No. 4,935,233 and U.S. Patent No. 4,751,180. The linker sequence may be from 1 to about 50 amino acids in length. Peptide sequences are not required when the first and second polypeptides have non-essential N-terminal amino acid regions that can be used to separate the functional domains and prevent steric interference.

The ligated polynucleotide sequences are operably linked to suitable transcriptional or translational regulatory elements. The regulatory elements responsible for expression of polynucleotides are located only 5' to the polynucleotide sequence encoding the first polypeptide. Similarly, stop codons required to end translation and transcription termination signals are only present 3' to the polynucleotide sequence encoding the second polypeptide.

Fusion proteins are also provided that comprise a polypeptide of the present invention together with an unrelated immunogenic protein. Preferably the immunogenic protein is capable of eliciting a recall response. Examples of such proteins include tetanus, tuberculosis and hepatitis proteins (see, for example, Stoute et al. *New Engl. J. Med.*, 336:86-91 (1997)).

Polypeptides of the present invention that comprise an immunogenic portion of a breast tumor protein may generally be used for immunotherapy of breast cancer, wherein the polypeptide stimulates the patient's own immune response to breast tumor cells. In further aspects, the present invention provides methods for using one or more of the immunoreactive polypeptides encoded by a polynucleotide

molecule having a sequence provided in SEQ ID NOS: 1- 97, 100 and 102-107 (or fusion proteins comprising one or more such polypeptides and/or polynucleotides encoding such polypeptides) for immunotherapy of breast cancer in a patient. As used herein, a "patient" refers to any warm-blooded animal, preferably a human. A patient may be afflicted with a disease, or may be free of detectable disease. Accordingly, the above immunoreactive polypeptides (or fusion proteins or polynucleotide molecules encoding such polypeptides) may be used to treat breast cancer or to inhibit the development of breast cancer. In a preferred embodiment, the polypeptides are administered either prior to or following surgical removal of primary tumors and/or treatment by administration of radiotherapy and conventional chemotherapeutic drugs.

In these aspects, the polypeptide or fusion protein is generally present within a pharmaceutical composition and/or a vaccine. Pharmaceutical compositions may comprise one or more polypeptides, each of which may contain one or more of the above sequences (or variants thereof), and a physiologically acceptable carrier. The vaccines may comprise one or more of such polypeptides and a non-specific immune response enhancer, wherein the non-specific immune response enhancer is capable of eliciting or enhancing an immune response to an exogenous antigen. Examples of non-specific-immune response enhancers include adjuvants, biodegradable microspheres (*e.g.*, polylactic galactide) and liposomes (into which the polypeptide is incorporated). Pharmaceutical compositions and vaccines may also contain other epitopes of breast tumor antigens, either incorporated into a combination polypeptide (*i.e.*, a single polypeptide that contains multiple epitopes) or present within a separate polypeptide.

Alternatively, a pharmaceutical composition or vaccine may contain polynucleotides encoding one or more of the above polypeptides, such that the polypeptide is generated *in situ*. In such pharmaceutical compositions and vaccines, the polynucleotide may be present within any of a variety of delivery systems known to those of ordinary skill in the art, including nucleic acid expression systems, bacteria and viral expression systems. Appropriate nucleic acid expression systems

contain the necessary polynucleotide sequences for expression in the patient (such as a suitable promoter). Bacterial delivery systems involve the administration of a bacterium (such as *Bacillus-Calmette-Guerrin*) that expresses an epitope of a breast tumor cell antigen on its cell surface. In a preferred embodiment, the polynucleotide molecules may be introduced using a viral expression system (e.g., vaccinia or other pox virus, retrovirus, or adenovirus), which may involve the use of a non-pathogenic (defective), replication competent virus. Suitable systems are disclosed, for example, in Fisher-Hoch et al., *PNAS* 86:317-321, 1989; Flexner et al., *Ann. N.Y. Acad. Sci.* 569:86-103, 1989; Flexner et al., *Vaccine* 8:17-21, 1990; U.S. Patent Nos. 4,603,112, 4,769,330, and 5,017,487; WO 89/01973; U.S. Patent No. 4,777,127; GB 2,200,651; EP 0,345,242; WO 91/02805; Berkner, *Biotechniques* 6:616-627, 1988; Rosenfeld et al., *Science* 252:431-434, 1991; Kolls et al., *PNAS* 91:215-219, 1994; Kass-Eisler et al., *PNAS* 90:11498-11502, 1993; Guzman et al., *Circulation* 88:2838-2848, 1993; and Guzman et al., *Cir. Res.* 73:1202-1207, 1993. Techniques for incorporating polynucleotides into such expression systems are well known to those of ordinary skill in the art.

The polynucleotides may also be "naked," as described, for example, in published PCT application WO 90/11092, and Ulmer et al., *Science* 259:1745-1749, 1993, reviewed by Cohen, *Science* 259:1691-1692, 1993. The uptake of naked polynucleotides may be increased by coating the polynucleotides onto biodegradable beads, which are efficiently transported into the cells.

Routes and frequency of administration, as well as dosage, will vary from individual to individual and may parallel those currently being used in immunotherapy of other diseases. In general, the pharmaceutical compositions and vaccines may be administered by injection (e.g., intracutaneous, intramuscular, intravenous or subcutaneous), intranasally (e.g., by aspiration) or orally. Between 1 and 10 doses may be administered over a 3-24 week period. Preferably, 4 doses are administered, at an interval of 3 months, and booster administrations may be given periodically thereafter. Alternate protocols may be appropriate for individual patients. A suitable dose is an amount of polypeptide or polynucleotide that is

effective to raise an immune response (cellular and/or humoral) against breast tumor cells in a treated patient. A suitable immune response is at least 10-50% above the basal (*i.e.*, untreated) level. In general, the amount of polypeptide present in a dose (or produced *in situ* by the polynucleotide in a dose) ranges from about 1 pg to about 100 mg per kg of host, typically from about 10 pg to about 1 mg, and preferably from about 100 pg to about 1 µg. Suitable dose sizes will vary with the size of the patient, but will typically range from about 0.01 mL to about 5 mL.

While any suitable carrier known to those of ordinary skill in the art may be employed in the pharmaceutical compositions of this invention, the type of carrier will vary depending on the mode of administration. For parenteral administration, such as subcutaneous injection, the carrier preferably comprises water, saline, alcohol, a lipid, a wax and/or a buffer. For oral administration, any of the above carriers or a solid carrier, such as mannitol, lactose, starch, magnesium stearate, sodium saccharine, talcum, cellulose, glucose, sucrose, and/or magnesium carbonate, may be employed. Biodegradable microspheres (*e.g.*, polylactic glycolide) may also be employed as carriers for the pharmaceutical compositions of this invention. Suitable biodegradable microspheres are disclosed, for example, in U.S. Patent Nos. 4,897,268 and 5,075,109.

Any of a variety of non-specific immune response enhancers may be employed in the vaccines of this invention. For example, an adjuvant may be included. Most adjuvants contain a substance designed to protect the antigen from rapid catabolism, such as aluminum hydroxide or mineral oil, and a nonspecific stimulator of immune response, such as lipid A, *Bordella pertussis* or *Mycobacterium tuberculosis*. Such adjuvants are commercially available as, for example, Freund's Incomplete Adjuvant and Complete Adjuvant (Difco Laboratories, Detroit, MI) and Merck Adjuvant 65 (Merck and Company, Inc., Rahway, NJ).

Polypeptides disclosed herein may also be employed in adoptive immunotherapy for the treatment of cancer. Adoptive immunotherapy may be broadly classified into either active or passive immunotherapy. In active immunotherapy, treatment relies on the *in vivo* stimulation of the endogenous host

immune system to react against tumors with the administration of immune response-modifying agents (for example, tumor vaccines, bacterial adjuvants, and/or cytokines).

In passive immunotherapy, treatment involves the delivery of biologic
5 reagents with established tumor-immune reactivity (such as effector cells or antibodies) that can directly or indirectly mediate antitumor effects and does not necessarily depend on an intact host immune system. Examples of effector cells include T lymphocytes (for example, CD8+ cytotoxic T-lymphocyte, CD4+ T-helper, gamma/delta T lymphocytes, tumor-infiltrating lymphocytes), killer cells (such as
10 Natural Killer cells, lymphokine-activated killer cells), B cells, or antigen presenting cells (such as dendritic cells and macrophages) expressing the disclosed antigens. The polypeptides disclosed herein may also be used to generate antibodies or anti-idiotypic antibodies (as in U.S. Patent No. 4,918,164), for passive immunotherapy.

The predominant method of procuring adequate numbers of T-cells for
15 adoptive immunotherapy is to grow immune T-cells *in vitro*. Culture conditions for expanding single antigen-specific T-cells to several billion in number with retention of antigen recognition *in vivo* are well known in the art. These *in vitro* culture conditions typically utilize intermittent stimulation with antigen, often in the presence of cytokines, such as IL-2, and non-dividing feeder cells. As noted above,
20 the immunoreactive polypeptides described herein may be used to rapidly expand antigen-specific T cell cultures in order to generate sufficient number of cells for immunotherapy. In particular, antigen-presenting cells, such as dendritic, macrophage, monocyte, fibroblast or B-cells, may be pulsed with immunoreactive polypeptides or polynucleotide sequence(s) may be introduced into antigen presenting
25 cells, using standard techniques well known in the art. For example, antigen presenting cells may be transfected or transduced with a polynucleotide sequence, wherein said sequence contains a promoter region appropriate for inducing expression, and can be expressed as part of a recombinant virus or other expression system. Several viral vectors may be used to transduce an antigen presenting cell,
30 including pox virus, vaccinia virus, and adenovirus. Antigen presenting cells may be

transfected with polynucleotide sequences disclosed herein by a variety of means, including gene-gun technology, lipid-mediated delivery, electroporation, osmotic shock, and particulate delivery mechanisms, resulting in efficient and acceptable expression levels as determined by one of ordinary skill in the art. For cultured T-cells to be effective in therapy, the cultured T-cells must be able to grow and distribute widely and to survive long term *in vivo*. Studies have demonstrated that cultured T-cells can be induced to grow *in vivo* and to survive long term in substantial numbers by repeated stimulation with antigen supplemented with IL-2 (see, for example, Cheever, M., et al, "Therapy With Cultured T Cells: Principles Revisited," *Immunological Reviews*, 157:177, 1997).

The polypeptides disclosed herein may also be employed to generate and/or isolate tumor-reactive T-cells, which can then be administered to the patient. In one technique, antigen-specific T-cell lines may be generated by *in vivo* immunization with short peptides corresponding to immunogenic portions of the disclosed polypeptides. The resulting antigen specific CD8+ CTL clones may be isolated from the patient, expanded using standard tissue culture techniques, and returned to the patient.

Alternatively, peptides corresponding to immunogenic portions of the polypeptides may be employed to generate tumor reactive T cell subsets by selective *in vitro* stimulation and expansion of autologous T cells to provide antigen-specific T cells which may be subsequently transferred to the patient as described, for example, by Chang et al. (*Crit. Rev. Oncol. Hematol.*, 22(3), 213, 1996). Cells of the immune system, such as T cells, may be isolated from the peripheral blood of a patient, using a commercially available cell separation system. The separated cells are stimulated with one or more of the immunoreactive polypeptides contained within a delivery vehicle, such as a microsphere, to provide antigen-specific T cells. The population of tumor antigen-specific T cells is then expanded using standard techniques and the cells are administered back to the patient.

In other embodiments, T-cell and/or antibody receptors specific for the polypeptides can be cloned, expanded, and transferred into other vectors or effector

cells for use in adoptive immunotherapy. In particular, T cells may be transfected with the appropriate genes to express the variable domains from tumor specific monoclonal antibodies as the extracellular recognition elements and joined to the T cell receptor signaling chains, resulting in T cell activation, specific lysis, and cytokine release. This enables the T cell to redirect its specificity in an MHC-independent manner. See for example, Eshhar, Z., *Cancer Immunol Immunother*, 45(3-4):131-6, 1997 and Hwu, P., et al, *Cancer Res*, 55(15):3369-73, 1995. Another embodiment may include the transfection of tumor antigen specific alpha and beta T cell receptor chains into alternate T cells, as in Cole, DJ, et al, *Cancer Res*, 55(4):748-52, 1995.

In further embodiments, syngeneic or autologous dendritic cells may be pulsed with peptides corresponding to at least an immunogenic portion of a polypeptide disclosed herein. The resulting antigen-specific dendritic cells may either be transferred into a patient, or employed to stimulate T cells to provide antigen-specific T cells which may, in turn, be administered to a patient. The use of peptide-pulsed dendritic cells to generate antigen-specific T cells and the subsequent use of such antigen-specific T cells to eradicate tumors in a murine model has been demonstrated by Cheever et al. (*Immunological Reviews*, 157:177, 1997).

Additionally, vectors expressing the disclosed polynucleotides may be introduced into stem cells taken from the patient and clonally propagated *in vitro* for autologous transplant back into the same patient.

In one specific embodiment, cells of the immune system, such as T cells, may be isolated from the peripheral blood of a patient, using a commercially available cell separation system, such as CellPro Incorporated's (Bothell, WA) CEPRATE™ system (see U.S. Patent No. 5,240,856; U.S. Patent No. 5,215,926; WO 89/06280; WO 91/16116 and WO 92/07243). The separated cells are stimulated with one or more of the immunoreactive polypeptides contained within a delivery vehicle, such as a microsphere, to provide antigen-specific T cells. The population of tumor antigen-specific T cells is then expanded using standard techniques and the cells are administered back to the patient.

Polypeptides of the present invention may also, or alternatively, be used to generate binding agents, such as antibodies or fragments thereof, that are capable of detecting metastatic human breast tumors. Binding agents of the present invention may generally be prepared using methods known to those of ordinary skill in the art, including the representative procedures described herein. Binding agents are capable of differentiating between patients with and without breast cancer, using the representative assays described herein. In other words, antibodies or other binding agents raised against a breast tumor protein, or a suitable portion thereof, will generate a signal indicating the presence of primary or metastatic breast cancer in at least about 20% of patients afflicted with the disease, and will generate a negative signal indicating the absence of the disease in at least about 90% of individuals without primary or metastatic breast cancer. Suitable portions of such breast tumor proteins are portions that are able to generate a binding agent that indicates the presence of primary or metastatic breast cancer in substantially all (*i.e.*, at least about 80%, and preferably at least about 90%) of the patients for which breast cancer would be indicated using the full length protein, and that indicate the absence of breast cancer in substantially all of those samples that would be negative when tested with full length protein. The representative assays described below, such as the two-antibody sandwich assay, may generally be employed for evaluating the ability of a binding agent to detect metastatic human breast tumors.

The ability of a polypeptide prepared as described herein to generate antibodies capable of detecting primary or metastatic human breast tumors may generally be evaluated by raising one or more antibodies against the polypeptide (using, for example, a representative method described herein) and determining the ability of such antibodies to detect such tumors in patients. This determination may be made by assaying biological samples from patients with and without primary or metastatic breast cancer for the presence of a polypeptide that binds to the generated antibodies. Such test assays may be performed, for example, using a representative procedure described below. Polypeptides that generate antibodies capable of detecting at least 20% of primary or metastatic breast tumors by such procedures are

considered to be useful in assays for detecting primary or metastatic human breast tumors. Polypeptide specific antibodies may be used alone or in combination to improve sensitivity.

Polypeptides capable of detecting primary or metastatic human breast
5 tumors may be used as markers for diagnosing breast cancer or for monitoring disease progression in patients. In one embodiment, breast cancer in a patient may be diagnosed by evaluating a biological sample obtained from the patient for the level of one or more of the above polypeptides, relative to a predetermined cut-off value. As used herein, suitable "biological samples" include blood, sera and urine.

10 The level of one or more of the above polypeptides may be evaluated using any binding agent specific for the polypeptide(s). A "binding agent," in the context of this invention, is any agent (such as a compound or a cell) that binds to a polypeptide as described above. As used herein, "binding" refers to a noncovalent association between two separate molecules (each of which may be free (*i.e.*, in
15 solution) or present on the surface of a cell or a solid support), such that a "complex" is formed. Such a complex may be free or immobilized (either covalently or noncovalently) on a support material. The ability to bind may generally be evaluated by determining a binding constant for the formation of the complex. The binding constant is the value obtained when the concentration of the complex is divided by
20 the product of the component concentrations. In general, two compounds are said to "bind" in the context of the present invention when the binding constant for complex formation exceeds about 10^3 L/mol. The binding constant may be determined using methods well known to those of ordinary skill in the art.

Any agent that satisfies the above requirements may be a binding
25 agent. For example, a binding agent may be a ribosome with or without a peptide component, an RNA molecule or a peptide. In a preferred embodiment, the binding partner is an antibody, or a fragment thereof. Such antibodies may be polyclonal, or monoclonal. In addition, the antibodies may be single chain, chimeric, CDR-grafted or humanized. Antibodies may be prepared by the methods described herein and by
30 other methods well known to those of skill in the art.

There are a variety of assay formats known to those of ordinary skill in the art for using a binding partner to detect polypeptide markers in a sample. *See, e.g., Harlow and Lane, Antibodies: A Laboratory Manual, Cold Spring Harbor Laboratory, 1988.* In a preferred embodiment, the assay involves the use of binding partner immobilized on a solid support to bind to and remove the polypeptide from the remainder of the sample. The bound polypeptide may then be detected using a second binding partner that contains a reporter group. Suitable second binding partners include antibodies that bind to the binding partner/polypeptide complex. Alternatively, a competitive assay may be utilized, in which a polypeptide is labeled with a reporter group and allowed to bind to the immobilized binding partner after incubation of the binding partner with the sample. The extent to which components of the sample inhibit the binding of the labeled polypeptide to the binding partner is indicative of the reactivity of the sample with the immobilized binding partner.

The solid support may be any material known to those of ordinary skill in the art to which the antigen may be attached. For example, the solid support may be a test well in a microtiter plate or a nitrocellulose or other suitable membrane. Alternatively, the support may be a bead or disc, such as glass, fiberglass, latex or a plastic material such as polystyrene or polyvinylchloride. The support may also be a magnetic particle or a fiber optic sensor, such as those disclosed, for example, in U.S. Patent No. 5,359,681. The binding agent may be immobilized on the solid support using a variety of techniques known to those of skill in the art, which are amply described in the patent and scientific literature. In the context of the present invention, the term "immobilization" refers to both noncovalent association, such as adsorption, and covalent attachment (which may be a direct linkage between the antigen and functional groups on the support or may be a linkage by way of a cross-linking agent). Immobilization by adsorption to a well in a microtiter plate or to a membrane is preferred. In such cases, adsorption may be achieved by contacting the binding agent, in a suitable buffer, with the solid support for a suitable amount of time. The contact time varies with temperature, but is typically between about 1 hour and about 1 day. In general, contacting a well of a plastic microtiter plate (such as

polystyrene or polyvinylchloride) with an amount of binding agent ranging from about 10 ng to about 10 μ g, and preferably about 100 ng to about 1 μ g, is sufficient to immobilize an adequate amount of binding agent.

Covalent attachment of binding agent to a solid support may generally
5 be achieved by first reacting the support with a bifunctional reagent that will react with both the support and a functional group, such as a hydroxyl or amino group, on the binding agent. For example, the binding agent may be covalently attached to supports having an appropriate polymer coating using benzoquinone or by
10 condensation of an aldehyde group on the support with an amine and an active hydrogen on the binding partner (*see, e.g.*, Pierce Immunotechnology Catalog and Handbook, 1991, at A12-A13).

In certain embodiments, the assay is a two-antibody sandwich assay. This assay may be performed by first contacting an antibody that has been immobilized on a solid support, commonly the well of a microtiter plate, with the
15 sample, such that polypeptides within the sample are allowed to bind to the immobilized antibody. Unbound sample is then removed from the immobilized polypeptide-antibody complexes and a second antibody (containing a reporter group) capable of binding to a different site on the polypeptide is added. The amount of second antibody that remains bound to the solid support is then determined using a
20 method appropriate for the specific reporter group.

More specifically, once the antibody is immobilized on the support as described above, the remaining protein binding sites on the support are typically blocked. Any suitable blocking agent known to those of ordinary skill in the art, such as bovine serum albumin or Tween 20TM (Sigma Chemical Co., St. Louis, MO). The
25 immobilized antibody is then incubated with the sample, and polypeptide is allowed to bind to the antibody. The sample may be diluted with a suitable diluent, such as phosphate-buffered saline (PBS) prior to incubation. In general, an appropriate contact time (*i.e.*, incubation time) is that period of time that is sufficient to detect the presence of polypeptide within a sample obtained from an individual with breast
30 cancer. Preferably, the contact time is sufficient to achieve a level of binding that is

at least about 95% of that achieved at equilibrium between bound and unbound polypeptide. Those of ordinary skill in the art will recognize that the time necessary to achieve equilibrium may be readily determined by assaying the level of binding that occurs over a period of time. At room temperature, an incubation time of about
5 30 minutes is generally sufficient.

Unbound sample may then be removed by washing the solid support with an appropriate buffer, such as PBS containing 0.1% Tween 20™. The second antibody, which contains a reporter group, may then be added to the solid support. Preferred reporter groups include enzymes (such as horseradish peroxidase),
10 substrates, cofactors, inhibitors, dyes, radionuclides, luminescent groups, fluorescent groups and biotin. The conjugation of antibody to reporter group may be achieved using standard methods known to those of ordinary skill in the art.

The second antibody is then incubated with the immobilized antibody-polypeptide complex for an amount of time sufficient to detect the bound
15 polypeptide. An appropriate amount of time may generally be determined by assaying the level of binding that occurs over a period of time. Unbound second antibody is then removed and bound second antibody is detected using the reporter group. The method employed for detecting the reporter group depends upon the nature of the reporter group. For radioactive groups, scintillation counting or
20 autoradiographic methods are generally appropriate. Spectroscopic methods may be used to detect dyes, luminescent groups and fluorescent groups. Biotin may be detected using avidin, coupled to a different reporter group (commonly a radioactive or fluorescent group or an enzyme). Enzyme reporter groups may generally be detected by the addition of substrate (generally for a specific period of time),
25 followed by spectroscopic or other analysis of the reaction products.

To determine the presence or absence of breast cancer, the signal detected from the reporter group that remains bound to the solid support is generally compared to a signal that corresponds to a predetermined cut-off value. In one preferred embodiment, the cut-off value is the average mean signal obtained when the
30 immobilized antibody is incubated with samples from patients without breast cancer.

In general, a sample generating a signal that is three standard deviations above the predetermined cut-off value is considered positive for breast cancer. In an alternate preferred embodiment, the cut-off value is determined using a Receiver Operator Curve, according to the method of Sackett et al., *Clinical Epidemiology: A Basic Science for Clinical Medicine*, Little Brown and Co., 1985, p. 106-7. Briefly, in this embodiment, the cut-off value may be determined from a plot of pairs of true positive rates (*i.e.*, sensitivity) and false positive rates (100%-specificity) that correspond to each possible cut-off value for the diagnostic test result. The cut-off value on the plot that is the closest to the upper left-hand corner (*i.e.*, the value that encloses the largest area) is the most accurate cut-off value, and a sample generating a signal that is higher than the cut-off value determined by this method may be considered positive. Alternatively, the cut-off value may be shifted to the left along the plot, to minimize the false positive rate, or to the right, to minimize the false negative rate. In general, a sample generating a signal that is higher than the cut-off value determined by this method is considered positive for breast cancer.

In a related embodiment, the assay is performed in a flow-through or strip test format, wherein the antibody is immobilized on a membrane, such as nitrocellulose. In the flow-through test, polypeptides within the sample bind to the immobilized antibody as the sample passes through the membrane. A second, labeled antibody then binds to the antibody-polypeptide complex as a solution containing the second antibody flows through the membrane. The detection of bound second antibody may then be performed as described above. In the strip test format, one end of the membrane to which antibody is bound is immersed in a solution containing the sample. The sample migrates along the membrane through a region containing second antibody and to the area of immobilized antibody. Concentration of second antibody at the area of immobilized antibody indicates the presence of breast cancer. Typically, the concentration of second antibody at that site generates a pattern, such as a line, that can be read visually. The absence of such a pattern indicates a negative result. In general, the amount of antibody immobilized on the membrane is selected to generate a visually discernible pattern when the biological sample contains a level

of polypeptide that would be sufficient to generate a positive signal in the two-antibody sandwich assay, in the format discussed above. Preferably, the amount of antibody immobilized on the membrane ranges from about 25 ng to about 1 μ g, and more preferably from about 50 ng to about 500 ng. Such tests can typically be performed with a very small amount of biological sample.

Of course, numerous other assay protocols exist that are suitable for use with the antigens or antibodies of the present invention. The above descriptions are intended to be exemplary only.

In another embodiment, the above polypeptides may be used as markers for the progression of breast cancer. In this embodiment, assays as described above for the diagnosis of breast cancer may be performed over time, and the change in the level of reactive polypeptide(s) evaluated. For example, the assays may be performed every 24-72 hours for a period of 6 months to 1 year, and thereafter performed as needed. In general, breast cancer is progressing in those patients in whom the level of polypeptide detected by the binding agent increases over time. In contrast, breast cancer is not progressing when the level of reactive polypeptide either remains constant or decreases with time.

Antibodies for use in the above methods may be prepared by any of a variety of techniques known to those of ordinary skill in the art. See, e.g., Harlow and Lane, *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory, 1988. In one such technique, an immunogen comprising the antigenic polypeptide is initially injected into any of a wide variety of mammals (e.g., mice, rats, rabbits, sheep and goats). In this step, the polypeptides of this invention may serve as the immunogen without modification. Alternatively, particularly for relatively short polypeptides, a superior immune response may be elicited if the polypeptide is joined to a carrier protein, such as bovine serum albumin or keyhole limpet hemocyanin. The immunogen is injected into the animal host, preferably according to a predetermined schedule incorporating one or more booster immunizations, and the animals are bled periodically. Polyclonal antibodies specific for the polypeptide may

then be purified from such antisera by, for example, affinity chromatography using the polypeptide coupled to a suitable solid support.

Monoclonal antibodies specific for the antigenic polypeptide of interest may be prepared, for example, using the technique of Kohler and Milstein, *Eur. J. Immunol.* 6:511-519, 1976, and improvements thereto. Briefly, these methods involve the preparation of immortal cell lines capable of producing antibodies having the desired specificity (*i.e.*, reactivity with the polypeptide of interest). Such cell lines may be produced, for example, from spleen cells obtained from an animal immunized as described above. The spleen cells are then immortalized by, for example, fusion with a myeloma cell fusion partner, preferably one that is syngeneic with the immunized animal. A variety of fusion techniques may be employed. For example, the spleen cells and myeloma cells may be combined with a nonionic detergent for a few minutes and then plated at low density on a selective medium that supports the growth of hybrid cells, but not myeloma cells. A preferred selection technique uses HAT (hypoxanthine, aminopterin, thymidine) selection. After a sufficient time, usually about 1 to 2 weeks, colonies of hybrids are observed. Single colonies are selected and tested for binding activity against the polypeptide. Hybridomas having high reactivity and specificity are preferred.

Monoclonal antibodies may be isolated from the supernatants of growing hybridoma colonies. In addition, various techniques may be employed to enhance the yield, such as injection of the hybridoma cell line into the peritoneal cavity of a suitable vertebrate host, such as a mouse. Monoclonal antibodies may then be harvested from the ascites fluid or the blood. Contaminants may be removed from the antibodies by conventional techniques, such as chromatography, gel filtration, precipitation, and extraction. The polypeptides of this invention may be used in the purification process in, for example, an affinity chromatography step.

Monoclonal antibodies of the present invention may also be used as therapeutic reagents, to diminish or eliminate breast tumors. The antibodies may be used on their own (for instance, to inhibit metastases) or coupled to one or more therapeutic agents. Suitable agents in this regard include radionuclides,

differentiation inducers, drugs, toxins, and derivatives thereof. Preferred radionuclides include ^{90}Y , ^{123}I , ^{125}I , ^{131}I , ^{186}Re , ^{188}Re , ^{211}At , and ^{212}Bi . Preferred drugs include methotrexate, and pyrimidine and purine analogs. Preferred differentiation inducers include phorbol esters and butyric acid. Preferred toxins include ricin, abrin,
5 diphtheria toxin, cholera toxin, gelonin, *Pseudomonas* exotoxin, *Shigella* toxin, and pokeweed antiviral protein.

A therapeutic agent may be coupled (*e.g.*, covalently bonded) to a suitable monoclonal antibody either directly or indirectly (*e.g.*, via a linker group). A direct reaction between an agent and an antibody is possible when each possesses a
10 substituent capable of reacting with the other. For example, a nucleophilic group, such as an amino or sulfhydryl group, on one may be capable of reacting with a carbonyl-containing group, such as an anhydride or an acid halide, or with an alkyl group containing a good leaving group (*e.g.*, a halide) on the other.

Alternatively, it may be desirable to couple a therapeutic agent and an
15 antibody via a linker group. A linker group can function as a spacer to distance an antibody from an agent in order to avoid interference with binding capabilities. A linker group can also serve to increase the chemical reactivity of a substituent on an agent or an antibody, and thus increase the coupling efficiency. An increase in chemical reactivity may also facilitate the use of agents, or functional groups on
20 agents, which otherwise would not be possible.

It will be evident to those skilled in the art that a variety of bifunctional or polyfunctional reagents, both homo- and hetero-functional (such as those described in the catalog of the Pierce Chemical Co., Rockford, IL), may be employed as the linker group. Coupling may be effected, for example, through amino
25 groups, carboxyl groups, sulfhydryl groups or oxidized carbohydrate residues. There are numerous references describing such methodology, *e.g.*, U.S. Patent No. 4,671,958, to Rodwell et al.

Where a therapeutic agent is more potent when free from the antibody portion of the immunoconjugates of the present invention, it may be desirable to use a
30 linker group which is cleavable during or upon internalization into a cell. A number

of different cleavable linker groups have been described. The mechanisms for the intracellular release of an agent from these linker groups include cleavage by reduction of a disulfide bond (*e.g.*, U.S. Patent No. 4,489,710, to Spitler), by irradiation of a photolabile bond (*e.g.*, U.S. Patent No. 4,625,014, to Senter et al.), by
5 hydrolysis of derivatized amino acid side chains (*e.g.*, U.S. Patent No. 4,638,045, to Kohn et al.), by serum complement-mediated hydrolysis (*e.g.*, U.S. Patent No. 4,671,958, to Rodwell et al.), and acid-catalyzed hydrolysis (*e.g.*, U.S. Patent No. 4,569,789, to Blattler et al.).

It may be desirable to couple more than one agent to an antibody. In
10 one embodiment, multiple molecules of an agent are coupled to one antibody molecule. In another embodiment, more than one type of agent may be coupled to one antibody. Regardless of the particular embodiment, immunoconjugates with more than one agent may be prepared in a variety of ways. For example, more than one agent may be coupled directly to an antibody molecule, or linkers which provide
15 multiple sites for attachment can be used. Alternatively, a carrier can be used.

A carrier may bear the agents in a variety of ways, including covalent bonding either directly or via a linker group. Suitable carriers include proteins such as albumins (*e.g.*, U.S. Patent No. 4,507,234, to Kato et al.), peptides and polysaccharides such as aminodextran (*e.g.*, U.S. Patent No. 4,699,784, to Shih et al.).
20 A carrier may also bear an agent by noncovalent bonding or by encapsulation, such as within a liposome vesicle (*e.g.*, U.S. Patent Nos. 4,429,008 and 4,873,088). Carriers specific for radionuclide agents include radiohalogenated small molecules and chelating compounds. For example, U.S. Patent No. 4,735,792 discloses representative radiohalogenated small molecules and their synthesis. A radionuclide
25 chelate may be formed from chelating compounds that include those containing nitrogen and sulfur atoms as the donor atoms for binding the metal, or metal oxide, radionuclide. For example, U.S. Patent No. 4,673,562, to Davison et al. discloses representative chelating compounds and their synthesis.

A variety of routes of administration for the antibodies and
30 immunoconjugates may be used. Typically, administration will be intravenous,

intramuscular, subcutaneous or in the bed of a resected tumor. It will be evident that the precise dose of the antibody/immunoconjugate will vary depending upon the antibody used, the antigen density on the tumor, and the rate of clearance of the antibody.

5 Diagnostic reagents of the present invention may also comprise at least a portion of a polynucleotide disclosed herein. For example, at least two oligonucleotide primers may be employed in a polymerase chain reaction (PCR) based assay to amplify breast tumor-specific cDNA derived from a biological sample, wherein at least one of the oligonucleotide primers is specific for a polynucleotide
10 encoding a breast tumor protein of the present invention. The presence of the amplified cDNA is then detected using techniques well known in the art, such as gel electrophoresis. Similarly, oligonucleotide probes specific for a polynucleotide encoding a breast tumor protein of the present invention may be used in a hybridization assay to detect the presence of an inventive polypeptide in a
15 biological sample.

 As used herein, the term "oligonucleotide primer/probe specific for a polynucleotide" means an oligonucleotide sequence that has at least about 60%, preferably at least about 75% and more preferably at least about 90%, identity to the polynucleotide in question, or an oligonucleotide sequence that is anti-sense to a
20 sequence that has at least about 60%, preferably at least about 75% and more preferably at least about 90%, identity to the polynucleotide in question. Oligonucleotide primers and/or probes which may be usefully employed in the inventive diagnostic methods preferably have at least about 10-40 nucleotides. In a preferred embodiment, the oligonucleotide primers comprise at least about 10
25 contiguous nucleotides of a polynucleotide disclosed herein or that is anti-sense to a polynucleotide sequence disclosed herein. Preferably, oligonucleotide probes for use in the inventive diagnostic methods comprise at least about 15 contiguous oligonucleotides of a polynucleotide that encodes one of the polypeptides disclosed herein or that is anti-sense to a sequence that encodes one of the polypeptides
30 disclosed herein. Techniques for both PCR based assays and hybridization assays are

well known in the art (see, for example, Mullis *et al. Ibid*; Ehrlich, *Ibid*). Primers or probes may thus be used to detect breast tumor-specific sequences in biological samples, including blood, urine and/or breast tumor tissue.

The following Examples are offered by way of illustration and not by
5 way of limitation.

EXAMPLES

Example 1

10 ISOLATION AND CHARACTERIZATION OF BREAST TUMOR POLYPEPTIDES

This Example describes the isolation of breast tumor polypeptides from a breast tumor cDNA library.

15 A human breast tumor cDNA expression library was constructed from a pool of breast tumor poly A⁺ RNA from three patients using a Superscript Plasmid System for cDNA Synthesis and Plasmid Cloning kit (BRL Life Technologies, Gaithersburg, MD 20897) following the manufacturer's protocol. Specifically, breast tumor tissues were homogenized with polytron (Kinematica, Switzerland) and total
20 RNA was extracted using Trizol reagent (BRL Life Technologies) as directed by the manufacturer. The poly A⁺ RNA was then purified using a Qiagen oligotex spin column mRNA purification kit (Qiagen, Santa Clarita, CA 91355) according to the manufacturer's protocol. First-strand cDNA was synthesized using the NotI/Oligo-dT18 primer. Double-stranded cDNA was synthesized, ligated with EcoRI/BstX I
25 adaptors (Invitrogen, Carlsbad, CA) and digested with NotI. Following size fractionation with Chroma Spin-1000 columns (Clontech, Palo Alto, CA 94303), the cDNA was ligated into the EcoRI/NotI site of pCDNA3.1 (Invitrogen, Carlsbad, CA) and transformed into ElectroMax *E. coli* DH10B cells (BRL Life Technologies) by electroporation.

Using the same procedure, a normal human breast cDNA expression library was prepared from a pool of four normal breast tissue specimens. The cDNA libraries were characterized by determining the number of independent colonies, the percentage of clones that carried insert, the average insert size and by sequence analysis. The breast tumor library contained 1.14×10^7 independent colonies, with more than 90% of clones having a visible insert and the average insert size being 936 base pairs. The normal breast cDNA library contained 6×10^6 independent colonies, with 83% of clones having inserts and the average insert size being 1015 base pairs. Sequencing analysis showed both libraries to contain good complex cDNA clones that were synthesized from mRNA, with minimal rRNA and mitochondrial DNA contamination sequencing.

cDNA library subtraction was performed using the above breast tumor and normal breast cDNA libraries, as described by Hara *et al.* (*Blood*, 84:189-199, 1994) with some modifications. Specifically, a breast tumor-specific subtracted cDNA library was generated as follows. Normal breast cDNA library (70 μ g) was digested with EcoRI, NotI, and SfuI, followed by a filling-in reaction with DNA polymerase Klenow fragment. After phenol-chloroform extraction and ethanol precipitation, the DNA was dissolved in 100 μ l of H₂O, heat-denatured and mixed with 100 μ l (100 μ g) of Photoprobe biotin (Vector Laboratories, Burlingame, CA), the resulting mixture was irradiated with a 270 W sunlamp on ice for 20 minutes. Additional Photoprobe biotin (50 μ l) was added and the biotinylation reaction was repeated. After extraction with butanol five times, the DNA was ethanol-precipitated and dissolved in 23 μ l H₂O to form the driver DNA.

To form the tracer DNA, 10 μ g breast tumor cDNA library was digested with BamHI and XhoI, phenol chloroform extracted and passed through Chroma spin-400 columns (Clontech). Following ethanol precipitation, the tracer DNA was dissolved in 5 μ l H₂O. Tracer DNA was mixed with 15 μ l driver DNA and 20 μ l of 2 x hybridization buffer (1.5 M NaCl/10 mM EDTA/50 mM HEPES pH 7.5/0.2% sodium dodecyl sulfate), overlaid with mineral oil, and heat-denatured completely. The sample was immediately transferred into a 68 °C water bath and

incubated for 20 hours (long hybridization [LH]). The reaction mixture was then subjected to a streptavidin treatment followed by phenol/chloroform extraction. This process was repeated three more times. Subtracted DNA was precipitated, dissolved in 12 μ l H₂O, mixed with 8 μ l driver DNA and 20 μ l of 2 x hybridization buffer, and
5 subjected to a hybridization at 68 °C for 2 hours (short hybridization [SH]). After removal of biotinylated double-stranded DNA, subtracted cDNA was ligated into BamHI/XhoI site of chloramphenicol resistant pBCSK⁺ (Stratagene, La Jolla, CA 92037) and transformed into ElectroMax *E. coli* DH10B cells by electroporation to generate a breast tumor specific subtracted cDNA library.

10 To analyze the subtracted cDNA library, plasmid DNA was prepared from 100 independent clones, randomly picked from the subtracted breast tumor specific library and characterized by DNA sequencing with a Perkin Elmer/Applied Biosystems Division Automated Sequencer Model 373A (Foster City, CA). Thirty-eight distinct cDNA clones were found in the subtracted breast tumor-specific cDNA
15 library. The determined 3' cDNA sequences for 14 of these clones are provided in SEQ ID NO: 1-14, with the corresponding 5' cDNA sequences being provided in SEQ ID NO: 15-28, respectively. The determined one strand (5' or 3') cDNA sequences for the remaining clones are provided in SEQ ID NO: 29-52. Comparison of these cDNA sequences with known sequences in the gene bank using the EMBL
20 and GenBank databases (Release 97) revealed no significant homologies to the sequences provided in SEQ ID NO: 3, 10, 17, 24 and 45-52. The sequences provided in SEQ ID NO: 1, 2, 4-9, 11-16, 18-23, 25-41, 43 and 44 were found to show at least some degree of homology to known human genes. The sequence of SEQ ID NO: 42 was found to show some homology to a known yeast gene.

25 cDNA clones isolated in the breast subtraction described above were colony PCR amplified and their mRNA expression levels in breast tumor, normal breast and various other normal tissues were determined using microarray technology (Synteni, Fremont, CA). Briefly, the PCR amplification products were dotted onto slides in an array format, with each product occupying a unique location in the array.
30 mRNA was extracted from the tissue sample to be tested, reverse transcribed, and

fluorescent-labeled cDNA probes were generated. The microarrays were probed with the labeled cDNA probes, the slides scanned and fluorescence intensity was measured. This intensity correlates with the hybridization intensity.

Data was analyzed using GEMTOOLS Software. Twenty one distinct
5 cDNA clones were found to be over-expressed in breast tumor and expressed at low levels in all normal tissues tested. The determined partial cDNA sequences for these clones are provided in SEQ ID NO: 53-73. Comparison of the sequences of SEQ ID NO: 53, 54 and 68-71 with those in the gene bank as described above, revealed some homology to previously identified human genes. No significant homologies were
10 found to the sequences of SEQ ID NO: 55-67, 72 (referred to as JJ 9434) and 73 (referred to as B535S). In further studies, full length cDNA sequences were obtained for the clones 1016F8 (SEQ ID NO: 56; also referred to as B511S) and 1016D12 (SEQ ID NO: 61; also referred to as B532S), and an extended cDNA sequence was obtained for 1012H8 (SEQ ID NO: 64; also referred to as B533S). These cDNA
15 sequences are provided in SEQ ID NO: 95-97, respectively, with the corresponding predicted amino acid sequences for B511S and B532S being provided in SEQ ID NO: 98 and 99, respectively.

Analysis of the expression of B511S in breast tumor tissues and in a variety of normal tissues (skin, PBMC, intestine, breast, stomach, liver, kidney, fetal
20 tissue, adrenal gland, salivary gland, spinal cord, large intestine, small intestine, bone marrow, brain, heart, colon and pancreas) by microarray, northern analysis and real time PCR, demonstrated that B511S is over-expressed in breast tumors, and normal breast, skin and salivary gland, with expression being low or undetectable in all other tissues tested.

25 Analysis of the expression of B532S in breast tumor tissue and in a variety of normal tissues (breast, PBMC, esophagus, HMEC, spinal cord, bone, thymus, brain, bladder, colon, liver, lung, skin, small intestine, stomach, skeletal muscle, pancreas, aorta, heart, spleen, kidney, salivary gland, bone marrow and adrenal gland) by microarray, Northern analysis and real time PCR, demonstrated that

B532S is over-expressed in 20-30% of breast tumors with expression being low or undetectable in all other tissues tested.

In a further experiment, cDNA fragments were obtained from two subtraction libraries derived by conventional subtraction, as described above and
5 analyzed by DNA microarray. In one instance the tester was derived from primary breast tumors, referred to as Breast Subtraction 2, or BS2. In the second instance, a metastatic breast tumor was employed as the tester, referred to as Breast Subtraction 3, or BS3. Drivers consisted of normal breast.

cDNA fragments from these two libraries were submitted as templates
10 for DNA microarray analysis, as described above. DNA chips were analyzed by hybridizing with fluorescent probes derived from mRNA from both tumor and normal tissues. Analysis of the data was accomplished by creating three groups from the sets of probes, referred to as breast tumor/mets, normal non-breast tissues, and metastatic breast tumors. Two comparisons were performed using the modified Gemtools
15 analysis. The first comparison was to identify templates with elevated expression in breast tumors. The second was to identify templates not recovered in the first comparison that yielded elevated expression in metastatic breast tumors. An arbitrary level of increased expression (mean of tumor expression versus the mean of normal tissue expression) was set at approximately 2.2.

20 In the first round of comparison to identify over-expression in breast tumors, two novel gene sequences were identified, hereinafter referred to as B534S and B538S (SEQ ID NO: 89 and 90, respectively), together with six sequences that showed some degree of homology to previously identified genes (SEQ ID NO: 74-79). The sequences of SEQ ID NO: 75 and 76 were subsequently determined to be
25 portions of B535S (SEQ ID NO: 73). In a second comparison to identify elevated expression in metastatic breast tumors, five novel sequences were identified, hereinafter referred to as B535S, B542S, B543S, P501S and B541S (SEQ ID NO: 73 and 91-94, respectively), as well as nine gene sequences that showed some homology to known genes (SEQ ID NO: 80-88). Clone B534S and B538S (SEQ ID NO: 89 and

90) were shown to be over-expressed in both breast tumors and metastatic breast tumors.

In a subsequent series of studies, 457 clones from Breast Subtraction 2 were analyzed by microarray on Breast Chip 3. As described above, a first comparison to identify over-expression in breast tumors over normal non-breast tissues was performed. This analysis yielded six cDNA clones that demonstrated elevated expression in breast tumor over normal non-breast tissues. Two of these clones, referred to as 1017C2 (SEQ ID NO: 102) and B546S (SEQ ID NO: 107) do not share significant homology to any known genes. Clone B511S also showed over-expression in breast tumor, which was previously described as 1016F8, with the determined cDNA sequence provided in SEQ ID NO: 95 and the predicted amino acid sequence provided in SEQ ID NO: 98. The remaining four clones over-expressed in breast tumor were found to share some degree of homology to Tumor Expression Enhanced Gene (SEQ ID NO: 103 and 104) Stromelysin-3 (SEQ ID NO: 105) or Collagen (SEQ ID NO: 106).

In the second comparison to determine genes with elevated expression in metastatic breast tumors over non-breast normal tissues, a profile similar to the first comparison was derived. The two putatively novel clones, 1017C2 and B546S, SEQ ID NO: 102 and 107, respectively, were overexpressed in metastatic breast tumors. In addition, Tumor Expression Enhanced Gene and B511S also showed elevated expression in metastatic breast tumors.

As described in U.S. Patent Application No. 08/806,099, filed February 25, 1997, the antigen P501S was isolated by subtracting a prostate tumor cDNA library with a normal pancreas cDNA library and with three genes found to be abundant in a previously subtracted prostate tumor specific cDNA library: human glandular kallikrein, prostate specific antigen (PSA), and mitochondria cytochrome C oxidase subunit II. The determined full-length cDNA sequence for P501S is provided in SEQ ID NO: 100, with the corresponding predicted amino acid sequence being provided in SEQ ID NO: 101. Expression of P501S in breast tumor was examined by microarray analysis. Over-expression was found in prostate tumor, breast tumor

and metastatic breast tumor, with negligible to low expression being seen in normal tissues. This data suggests that P501S may be over-expressed in various breast tumors as well as in prostate tumors.

5

Example 2

GENERATION OF HUMAN CD8+ CYTOTOXIC T-CELLS THAT RECOGNIZE ANTIGEN PRESENTING CELLS EXPRESSING BREAST TUMOR ANTIGENS

10 This Example illustrates the generation of T cells that recognize target cells expressing the antigen B511S, also known as 1016-F8 (SEQ ID NO: 56). Human CD8+ T cells were primed *in-vitro* to the B511S gene product using dendritic cells infected with a recombinant vaccinia virus engineered to express B511S as follows (also see Yee et al., Journal of Immunology (1996) 157 (9):4079-86).

15 Dendritic cells (DC) were generated from peripheral blood derived monocytes by differentiation for 5 days in the presence of 50 µg/ml GMCSF and 30 µg/ml IL-4. DC were harvested, plated in wells of a 24-well plate at a density of 2×10^5 cells/well and infected for 12 hours with B511S expressing vaccinia at a multiplicity of infection of 5. DC were then matured overnight by the addition of 3 µg/ml CD40-

20 Ligand and UV irradiated at 100µW for 10 minutes. CD8+ T cells were isolated using magnetic beads, and priming cultures were initiated in individual wells (typically in 24 wells of a 24-well plate) using 7×10^5 CD8+ T cells and 1×10^6 irradiated CD8-depleted PBMC. IL-7 at 10 ng/ml was added to cultures at day 1. Cultures were re-stimulated every 7-10 days using autologous primary fibroblasts

25 retrovirally transduced with B511S and the costimulatory molecule B7.1. Cultures were supplemented at day 1 with 15 I.U. of IL-2. Following 4 such stimulation cycles, CD8+ cultures were tested for their ability to specifically recognize autologous fibroblasts transduced with B511S using an interferon-γ Elispot assay (see Lalvani et al J. Experimental Medicine (1997) 186:859-965). Briefly, T cells from

30 individual microcultures were added to 96-well Elispot plates that contained autologous fibroblasts transduced to express either B511S or as a negative control

antigen EGFP, and incubated overnight at 37° C; wells also contained IL-12 at 10 ng/ml. Cultures were identified that specifically produced interferon- γ only in response to B511S transduced fibroblasts; such lines were further expanded and also cloned by limiting dilution on autologous B-LCL retrovirally transduced with B511S.

- 5 Lines and clones were identified that could specifically recognize autologous B-LCL transduced with B511S but not autologous B-LCL transduced with the control antigens EGFP or HLA-A3. An example demonstrating the ability of human CTL cell lines derived from such experiments to specifically recognize and lyse B511S expressing targets is presented in Figure 1.

10

Example 3

SYNTHESIS OF POLYPEPTIDES

- Polypeptides may be synthesized on an Perkin Elmer/Applied Biosystems Division 430A peptide synthesizer using Fmoc chemistry with HPTU (O-Benzotriazole-N,N,N',N'-tetramethyluronium hexafluorophosphate) activation. A Gly-Cys-Gly sequence may be attached to the amino terminus of the peptide to provide a method of conjugation, binding to an immobilized surface, or labeling of the peptide. Cleavage of the peptides from the solid support may be carried out using
- 20 the following cleavage mixture: trifluoroacetic acid:ethanedithiol:thioanisole:water:phenol (40:1:2:2:3). After cleaving for 2 hours, the peptides may be precipitated in cold methyl-t-butyl-ether. The peptide pellets may then be dissolved in water containing 0.1% trifluoroacetic acid (TFA) and lyophilized prior to purification by C18 reverse phase HPLC. A gradient of 0%-60%
- 25 acetonitrile (containing 0.1% TFA) in water (containing 0.1% TFA) may be used to elute the peptides. Following lyophilization of the pure fractions, the peptides may be characterized using electrospray or other types of mass spectrometry and by amino acid analysis.

From the foregoing, it will be appreciated that, although specific embodiments of the invention have been described herein for the purposes of illustration, various modifications may be made without deviating from the spirit and scope of the invention.

CLAIMS

1. An isolated polypeptide comprising an immunogenic portion of a breast protein or a variant thereof, wherein said protein comprises an amino acid sequence encoded by a polynucleotide comprising a sequence selected from the group consisting of: (a) nucleotide sequences recited in SEQ ID NOS: 3, 10, 17, 24, 45-52, 55-67, 72, 73, 89-97, 102 and 107; (b) complements of said nucleotide sequences; and (c) sequences that hybridize to a sequence of (a) or (b) under moderately stringent conditions.
2. The isolated polypeptide of claim 1, wherein the polypeptide comprises an amino acid sequence selected from the group consisting of SEQ ID NO: 98,99 and 101.
3. An isolated polynucleotide comprising a nucleotide sequence encoding the polypeptide of any one of claims 1 and 2.
4. An isolated polynucleotide comprising a sequence provided in SEQ ID NOS: 3, 10, 17, 24, 45-52, 55-67, 72, 73, 89-97, 102 and 107.
5. An expression vector comprising a polynucleotide according to any one of claims 3 and 4.
6. A host cell transformed with the expression vector of claim 5.
7. The host cell of claim 6 wherein the host cell is selected from the group consisting of *E. coli*, yeast and mammalian cell lines.
8. A pharmaceutical composition comprising the polypeptide of claim 1 and a physiologically acceptable carrier.
9. A vaccine comprising the polypeptide of claim 1 and a non-specific immune response enhancer.

10. The vaccine of claim 9 wherein the non-specific immune response enhancer is an adjuvant.

11. A vaccine comprising an isolated polynucleotide of any one of claims 3 and 4, and a non-specific immune response enhancer.

12. The vaccine of claim 11 wherein the non-specific immune response enhancer is an adjuvant.

13. A pharmaceutical composition for the treatment of breast cancer comprising a polypeptide and a physiologically acceptable carrier, the polypeptide comprising an immunogenic portion of a breast protein, wherein said protein comprises an amino acid sequence encoded by a polynucleotide comprising a sequence selected from the group consisting of: (a) nucleotide sequences recited in SEQ ID NOS: 1, 2, 4-9, 11-16, 18-23, 25-44, 53, 54, 68-71, 74-88 and 103-106; (b) complements of said nucleotide sequences; and (c) sequences that hybridize to a sequence of (a) or (b) under moderately stringent conditions.

14. A vaccine for the treatment of breast cancer comprising a polypeptide and a non-specific immune response enhancer, said polypeptide comprising an immunogenic portion of a breast protein, wherein said protein comprises an amino acid sequence encoded by a polynucleotide comprising a sequence selected from the group consisting of: (a) nucleotide sequences recited in SEQ ID NOS: 1, 2, 4-9, 11-16, 18-23, 25-44, 53, 54, 68-71, 74-88 and 103-106; (b) complements of said nucleotide sequences; and (c) sequences that hybridize to a sequence of (a) or (b) under moderately stringent conditions.

15. The vaccine of claim 14 wherein the non-specific immune response enhancer is an adjuvant.

16. A vaccine for the treatment of breast cancer comprising a polynucleotide and a non-specific immune response enhancer, the polynucleotide

comprising a sequence selected from the group consisting of: (a) nucleotide sequences recited in SEQ ID NOS: 1, 2, 4-9, 11-16, 18-23, 25-44, 53, 54, 68-71, 74-88 and 103-106; (b) complements of said nucleotide sequences; and (c) sequences that hybridize to a sequence of (a) or (b) under moderately stringent conditions.

17. The vaccine of claim 16, wherein the non-specific immune response enhancer is an adjuvant.

18. A pharmaceutical composition according to any one of claims 8 and 13, for use in the manufacture of a medicament for inhibiting the development of breast cancer in a patient.

19. A vaccine according to any one of claims 9, 11, 14 or 16, for use in the manufacture of a medicament for inhibiting the development of breast cancer in a patient.

20. A fusion protein comprising at least one polypeptide according to claim 1.

21. A pharmaceutical composition comprising a fusion protein according to claim 20 and a physiologically acceptable carrier.

22. A vaccine comprising a fusion protein according to claim 20 and a non-specific immune response enhancer.

23. The vaccine of claim 22 wherein the non-specific immune response enhancer is an adjuvant.

24. A pharmaceutical composition according to claim 21, for use in manufacture of a medicament for inhibiting the development of breast cancer in a patient.

25. A vaccine according to claim 22, for use in the manufacture of a medicament for inhibiting the development of breast cancer in a patient.

26. A method for detecting breast cancer in a patient, comprising:

(a) contacting a biological sample from a patient with a binding agent which is capable of binding to a polypeptide, the polypeptide comprising an immunogenic portion of a breast protein, wherein said protein comprises an amino acid sequence encoded by a polynucleotide comprising a sequence selected from the group consisting of nucleotide sequences recited in SEQ ID NOS: 1-97, 100 and 102-107, complements of said nucleotide sequences and sequences that hybridize to a sequence provided in SEQ ID NO: 1-97, 100 and 102-107 under moderately stringent conditions; and

(b) detecting in the sample a protein or polypeptide that binds to the binding agent, thereby detecting breast cancer in the patient.

27. The method of claim 26 wherein the binding agent is a monoclonal antibody.

28. The method of claim 27 wherein the binding agent is a polyclonal antibody.

29. A method for monitoring the progression of breast cancer in a patient, comprising:

(a) contacting a biological sample from a patient with a binding agent that is capable of binding to a polypeptide, said polypeptide comprising an immunogenic portion of a breast protein, wherein said protein comprises an amino acid sequence encoded by a polynucleotide comprising a sequence selected from the group consisting of nucleotide sequences recited in SEQ ID NOS: 1-97, 100 and 102-107, complements of said nucleotide sequences and sequences that hybridize to a sequence provided in SEQ ID NO: 1-97, 100 and 102-107 under moderately stringent conditions;

(b) determining in the sample an amount of a protein or polypeptide that binds to the binding agent;

(c) repeating steps (a) and (b); and
(d) comparing the amount of polypeptide detected in steps (b) and (c)
to monitor the progression of breast cancer in the patient.

30. A monoclonal antibody that binds to a polypeptide comprising an immunogenic portion of a breast protein or a variant of said protein that differs only in conservative substitutions and/or modifications, wherein said protein comprises an amino acid sequence encoded by a polynucleotide comprising a sequence selected from the group consisting of: (a) nucleotide sequences recited in SEQ ID NOS: 3, 10, 17, 24, 45-52, 55-67, 72, 73, 89-97, 102 and 107; (b) complements of said nucleotide sequences; and (c) sequences that hybridize to a sequence of (a) or (b) under moderately stringent conditions.

31. A monoclonal antibody according to claim 30, for use in the manufacture of a medicament for inhibiting the development of breast cancer in a patient.

32. The monoclonal antibody of claim 31 wherein the monoclonal antibody is conjugated to a therapeutic agent.

33. A method for detecting breast cancer in a patient comprising:

- (a) contacting a biological sample from a patient with at least two oligonucleotide primers in a polymerase chain reaction, wherein at least one of the oligonucleotides is specific for a polynucleotide encoding a polypeptide comprising an immunogenic portion of a breast protein, said protein comprising an amino acid sequence encoded by a polynucleotide comprising a sequence selected from the group consisting of nucleotide sequences recited in SEQ ID NO: 1-97, 100 and 102-107, complements of said nucleotide sequences and sequences that hybridize to a sequence of SEQ ID NO: 1-97, 100 and 102-107 under moderately stringent conditions; and
(b) detecting in the sample a polynucleotide sequence that amplifies in the presence of the oligonucleotide primers, thereby detecting breast cancer.

34. The method of claim 33, wherein at least one of the oligonucleotide primers comprises at least about 10 contiguous nucleotides of a polynucleotide comprising a sequence selected from SEQ ID NOS: 1-97, 100 and 102-107.

35. A diagnostic kit comprising:

- (a) one or more monoclonal antibodies of claim 30; and
- (b) a detection reagent.

36. A diagnostic kit comprising:

- (a) one or more monoclonal antibodies that bind to a polypeptide encoded by a polynucleotide comprising a nucleotide sequence selected from the group consisting of SEQ ID NOS: 1, 2, 4-9, 11-16, 18-23, 25-44, 53, 54, 68-71, 74-88 and 103-106, complements of said sequences, and sequences that hybridize to a sequence of SEQ ID NO: 1, 2, 4-9, 11-16, 18-23, 25-44, 53, 54, 68-71, 74-88 or 103-106 under moderately stringent conditions; and
- (b) a detection reagent.

37. The kit of claims 35 or 36 wherein the monoclonal antibodies are immobilized on a solid support.

38. The kit of claim 37 wherein the solid support comprises nitrocellulose, latex or a plastic material.

39. The kit of claims 35 or 36 wherein the detection reagent comprises a reporter group conjugated to a binding agent.

40. The kit of claim 39 wherein the binding agent is selected from the group consisting of anti-immunoglobulins, Protein G, Protein A and lectins.

41. The kit of claim 39 wherein the reporter group is selected from the group consisting of radioisotopes, fluorescent groups, luminescent groups, enzymes, biotin and dye particles.

42. A diagnostic kit comprising at least two oligonucleotide primers, at least one of the oligonucleotide primers being specific for a polynucleotide encoding a polypeptide comprising an immunogenic portion of a breast protein, said protein comprising an amino acid sequence encoded by a polynucleotide comprising a sequence selected from the group consisting of nucleotide sequences recited in SEQ ID NOS: 1-97, 100 and 102-107 complements of said nucleotide sequences and sequences that hybridize to a sequence of SEQ ID NO: 1-97, 100 and 102-107 under moderately stringent conditions.

43. A diagnostic kit of claim 42 wherein at least one of the oligonucleotide primers comprises at least about 10 contiguous nucleotides of a polynucleotide comprising a sequence selected from SEQ ID NOS: 1-97, 100 and 102-107.

44. A method for detecting breast cancer in a patient, comprising:

- (a) obtaining a biological sample from the patient;
- (b) contacting the biological sample with an oligonucleotide probe specific for a polynucleotide encoding a polypeptide comprising an immunogenic portion of a breast protein, said protein comprising an amino acid sequence encoded by a polynucleotide comprising a sequence selected from the group consisting of nucleotide sequences recited in SEQ ID NOS: 1-97, 100 and 102-107, complements of said nucleotide sequences and sequences that hybridize to a sequence of SEQ ID NO: 1-97, 100 and 102-107 under moderately stringent conditions; and
- (c) detecting in the sample a polynucleotide sequence that hybridizes to the oligonucleotide probe, thereby detecting breast cancer in the patient.

45. The method of claim 44 wherein the oligonucleotide probe comprises at least about 15 contiguous nucleotides of a polynucleotide comprising a sequence selected from the group consisting of SEQ ID NOS: 1-97, 100 and 102-107.

46. A diagnostic kit comprising an oligonucleotide probe specific for a polynucleotide encoding a polypeptide comprising an immunogenic portion of a breast protein, said protein comprising an amino acid sequence encoded by a polynucleotide comprising a sequence selected from the group consisting of nucleotide sequences recited in SEQ ID NOS: 1-97, 100 and 102-107, complements of said nucleotide sequences, and sequences that hybridize to a sequence of SEQ ID NO: 1-97, 100 and 102-107 under moderately stringent conditions.

47. The diagnostic kit of claim 46, wherein the oligonucleotide probe comprises at least about 15 contiguous nucleotides of a polynucleotide comprising a sequence selected from the group consisting of SEQ ID NOS: 1-97, 100 and 102-107.

48. A method for treating breast cancer in a patient, comprising the steps of:

- (a) obtaining peripheral blood cells from the patient;
- (b) incubating the cells in the presence of at least one polypeptide of any one of claims 1 and 2, such that T cells proliferate; and administering the proliferated T cells to the patient.

49. A method for treating breast cancer in a patient, comprising the steps of:

- (a) obtaining peripheral blood cells from the patient;
- (b) incubating the cells in the presence of at least one polynucleotide of any one of claims 3 and 4, such that T cells proliferate; and
- (c) administering the proliferated T cells to the patient.

50. The method of any one of claims 48 and 49 wherein the step of incubating the cells is repeated one or more times.

51. The method of any one of claims 48 and 49 wherein step (a) further comprises separating the T cells from the peripheral blood cells and the cells incubated in step (b) are the T cells.

52. The method of any one of claims 48 and 49 wherein step (a) further comprises separating CD4⁺ cells or CD8⁺ cells from the peripheral blood cells and the cells proliferated in step (b) are CD4⁺ or CD8⁺ T cells.

53. The method of any one of claims 48 and 49 wherein step (b) further comprises cloning at least one T cell that proliferated in the presence of the polypeptide.

54. A composition for the treatment of breast cancer in a patient, comprising T cells proliferated in the presence of a polypeptide of any one of claims 1 and 2, in combination with a pharmaceutically acceptable carrier.

55. A composition for the treatment of breast cancer in a patient comprising T cells proliferated in the presence of a polynucleotide of any one of claims 3 and 4, in combination with a pharmaceutically acceptable carrier.

56. A method for treating breast cancer in a patient, comprising the steps of:

- (a) incubating antigen presenting cells in the presence of at least one polypeptide of any one of claims 1 and 2; and
- (b) administering to the patient the incubated antigen presenting cells.

57. A method for treating breast cancer in a patient, comprising the steps of:

- (a) incubating antigen presenting cells in the presence of at least one polynucleotide of any one of claims 3 and 4; and
- (b) administering to the patient the incubated antigen presenting

cells.

58. The method of claims 56 or 57 wherein the antigen presenting cells are selected from the group consisting of dendritic cells, macrophage cells, monocyte cells, fibroblast cells, B-cells or combinations thereof.

59. A composition for the treatment of breast cancer in a patient, comprising antigen presenting cells incubated in the presence of a polypeptide of any one of claims 1 and 2, in combination with a pharmaceutically acceptable carrier.

60. A composition for the treatment of breast cancer in a patient, comprising antigen presenting cells incubated in the presence of a polynucleotide of any one of claims 3 and 4, in combination with a pharmaceutically acceptable carrier.




FIGURE 1A




FIGURE 1B

Figure 1: Specific lytic activity of B511s-specific CTL clones 3-6-8 and 3-6-7 measured on autologous LCL transduced with B511s (filled squares) or HLA-A3 (open squares). Each data point is the average of triplicate measurements.

FIGS. 1A AND 1B

SEQUENCE LISTING

<110> Corixa Corporation
 Reed, Steven G.
 Xu, Jiangchun
 Dillon, Davin C.

<120> COMPOUNDS FOR IMMUNOTHERAPY AND
 DIAGNOSIS OF BREAST CANCER AND METHODS FOR THEIR USE

<130> 210121.44602PC

<140> PCT

<141> 2000-04-10

<160> 107

<170> FastSEQ for Windows Version 3.0

<210> 1

<211> 402

<212> DNA

<213> Homo sapien

<220>

<221> misc_feature

<222> (1)...(402)

<223> n = A,T,C or G

<400> 1

tttttttttt tttttaggag aactgaatca aacagatttt attcaacttt ttagatgagg	60
aaaacaaatn atacgaaatn ngtcataaga aatgctttct tataccacta tctcaaacca	120
ctttcaatat tttaaaaaat gctcacgcag caaatatgaa aagctncaac acttcccttt	180
gttaacttgc tgcaatnaat gcaactttta canacatata aatttcttct gtatcttaaa	240
agttnaatta ctaattttta tgatnttnt caagatnttt attcatatac ttttaatgac	300
tcnttgccna tacatacnta ttttctttac ttttttttta cnatnggcca acagctttca	360
ngcagncnc aaaaatctta ccggttaatt acacggggtt gt	402

<210> 2

<211> 424

<212> DNA

<213> Homo sapien

<220>

<221> misc_feature

<222> (1)...(424)

<223> n = A,T,C or G

<400> 2

tttttttttt ttttttaaag gtacacattt ctttttcatt ctgtttnatg cagcaaataa	60
ttcgttggca tcttctctgt gatgggcagc ttgctaaaat tanactcagg ccccttagct	120
ncatttccaa cttagccac gctttcaacc nngccnaaca aagaaaatca gttngggtta	180
aattctttgc tgganacaaa gaactacatt cctttgtaaa tnatgctttg tttgctctgt	240

gcaaacncag attgaagggg anaagganac ttntggggac ggaaacaact ngnagaagca 300
gganccgccc agggncattt cctcaccatg cttaatcttg cnctcacttg cngggcacca 360
ttaaacttgg tgcaaaaggc gcaattggtg nanggaaccc cacaccttcc ttaaaaagca 420
gggc 424

<210> 3
<211> 421
<212> DNA
<213> Homo sapien

<220>
<221> misc_feature
<222> (1)...(421)
<223> n = A,T,C or G

<400> 3
tttttttttt tttttcccaa tttaaaaaag cttttttcat acttcaatta caccanactt 60
aatnatttca tgagtaaadc ngacattatt atttnaaaat ttgcataattt aaaatttgna 120
tcanttactt ccagactggt tgcanaatga agggaggatc actcaagngc tgatctcnca 180
ctntctgcag tctnctgtcc tgtgcccggg ctaatggatc gacactanat ggacagntcn 240
cagatcttcc gttcttntcc cttccccaat ttcnaccnc tccccttctt ncccggatcn 300
tttggggaca tgntaatatt gcnatcctta aaccctgccc gccanggggc ccnancctag 360
gggtgggttaa tgttcgneng gcttnttgac cncctgcgcc ctttnantcc naaccccaag 420
c 421

<210> 4
<211> 423
<212> DNA
<213> Homo sapien

<220>
<221> misc_feature
<222> (1)...(423)
<223> n = A,T,C or G

<400> 4
tttttttatt tttttttcta tttntnntat ttntgnggt tcctgtgtgt aattagnang 60
tgtgtatgcg tangtacnta tgtntgcata tttaacctgt tncctttcca tttttaaaat 120
aaaatctcaa natngtantt ggttnatggg agtaaanaga gactatngat naattttaac 180
atggacacng tgaaatgtag ccgctnatca nttaaaaact tcattttgaa ggccttttnc 240
cctccnaata aaaatnccng gccctactgg gtttaagcaac attgcatntc taaagaaacc 300
acatgcanac nagttaaacc tgtgnactgg tcangcaaac cnanntggaa nanaagggnn 360
ttcncccan ggacantcng aattttttta acaaattacn atnccccccc ngggggagcc 420
tgt 423

<210> 5
<211> 355
<212> DNA
<213> Homo sapien

<220>
<221> misc_feature
<222> (1)...(355)
<223> n = A,T,C or G

```

<400> 5
acgaccacct natttcgtat ctttcaactc ttttcgaccg gacctcttat tcggaagcgt      60
tccaggaaga caggtctcaa cttagggatc agatcacgtt atcaacgctc tgggatcgct      120
gcaacctggc acttcaagga agtgcaccga tnacgtctag accggccaac acagatctag      180
aggtggccaa ctgatcactg taggagctga ctggcaanan tcaaccgggc cccaaccnag      240
agtgaccaan acnaccattn aggatcacc cagggcactc ctctccttag ggccaaccna      300
ccaaacggct ggccaatggg ggggtttaat atttggttna aaaattgatt ttaaa          355

```

```

<210> 6
<211> 423
<212> DNA
<213> Homo sapien

```

```

<220>
<221> misc_feature
<222> (1)...(423)
<223> n = A,T,C or G

```

```

<400> 6
tttttttttt tttttggaca ggaagtaaaa tttattggtn antattaana ggggggcagc      60
acattggaag ccctcatgan tgcagggccc gccacttgtc cagagggcca cnattgggga      120
tgtacttaac cccacagccn tctgggatna gccgcttttc agccaccatn tcttcaaatt      180
catcagcatt aaacttggtg aanccccact tctttaagat ntgnatcttc tggcgggcag      240
naaacttgaa cttggccctg cgagggcct caatcacatg ctcttggttc tgcagcttg      300
tgcgnaagga cntaatnact tggccnatgt gaaccctggc cacantgccc tggggctttc      360
caaaggcacc tcgcaagcct ntttggancc tgnccgcccc ngcacaggga caacatcttg      420
ttt                                          423

```

```

<210> 7
<211> 410
<212> DNA
<213> Homo sapien

```

```

<220>
<221> misc_feature
<222> (1)...(410)
<223> n = A,T,C or G

```

```

<400> 7
ttcgcaactg ctaaaacaaa ccgccttgca aagttnghaa aatttatcaa tggaccaa      60
aatgctcata tccnacaagt tggtagccgt tnttatnata aaaaaatgta tnatgctcct      120
nanttggtgt acaataatgt tccaatttng gacnttcggc atctaccctg gttcacctgg      180
gtaaatatca ggcagctttt gatgggggcta ggaaagctaa cagtactcga acatgggaaa      240
gaggtctgct tcgccngtgt anatgggaaa naattccgtc ttgctcngat ttgtggactt      300
catattgttg tacatgcaga tgaatnngaa gaacttgtea actactatca ggatcgtggc      360
tttttnnaaa agctnatcac catgttgga gcggcactng gacttgagcg          410

```

```

<210> 8
<211> 274
<212> DNA
<213> Homo sapien

```

```

<220>
<221> misc_feature
<222> (1)...(274)

```

<223> n = A,T,C or G

<400> 8

tttttttttt	tttttaggtc	atacatattt	tttattataa	canatatntg	tatatacata	60
taatatatgt	gtatatatcc	acgtgtgtgt	gtgtgtatca	aaaacaacan	aanttttagtg	120
atctatatct	ntngctcaca	tatgcatggg	agataccagt	aaaaaataag	tnaatctcca	180
taatatgttt	taaaactcan	anaaatcnga	gagactnaaa	gaaaacgtn	atcannatga	240
ttgtngataa	tcttgaanaa	tnacnaaaac	atat			274

<210> 9

<211> 322

<212> DNA

<213> Homo sapien

<220>

<221> misc_feature

<222> (1)...(322)

<223> n = A,T,C or G

<400> 9

tttttttttt	ttttgtgcct	tattgcaccg	gcnanaactt	ctagcactat	attaaactca	60
ataagagtga	taagtgtgaa	aatccttgcc	ttctctttaa	tcttaatgna	naggcatctg	120
gtttttcacc	attaantgta	ataatggctn	tatgtatttt	tatnnatggg	cttnatggag	180
ttaaaaaagt	tttctctnt	ccctngttat	ctaanagttt	tnatcaaaaa	tgggtataat	240
atttngttca	gtacttttnc	ctgcacctat	agatatgatn	ctgttatttt	ttcttcttng	300
cctnnanata	tgatggatna	ca				322

<210> 10

<211> 425

<212> DNA

<213> Homo sapien

<220>

<221> misc_feature

<222> (1)...(425)

<223> n = A,T,C or G

<400> 10

tttttttttt	tttttattct	gcagccatta	aatgctgaac	actagatnct	tatttgtgga	60
ggtcacaaaa	taagtacaga	atatnacaca	cgccctgccc	ataaaaagca	cagctcccag	120
ttctatatatt	acaatatctc	tggaattcca	ccttcccttc	taatttgact	aatatttctg	180
cttctcaggc	agcagcgctt	tctggcaacc	ataagaacca	acntgnggac	taggtcgggtg	240
ggccaaggat	caggaaacag	aanaatggaa	gnagccccc	tgacnctatt	aancntnaa	300
actatctnaa	ctgctagtgt	tcaggcttta	aatcatgtaa	natacgtgtc	cttnttgctg	360
caaccggaag	catcctagat	ggtagactct	ctccagggtc	caggaaaaga	tcccaaattg	420
caggn						425

<210> 11

<211> 424

<212> DNA

<213> Homo sapien

<220>

<221> misc_feature

<222> (1)...(424)

<223> n = A,T,C or G

<400> 11

ttttnttant	ttttttancc	nctnntccnn	tntgttnag	ggggtacca	atttctttat	60
ttaaagggaat	ggtacaaatc	aaaaaactta	atttaatttt	tnggtacaac	ttatagaaaa	
120ggttaaggaa	acccaacat	gcatgcactg	ccttggtaac	cagggnatc	ccccncggct	
180						
ntgggggaaat	tagcccaang	ctnagctttc	attatcactn	tccccaggg	tntgcttttc	240
aaaaaaattt	nccgccnagc	cnaatccggg	cncctccatc	tggcgcaant	tggtcacttg	300
gtcccccnat	tctttaangg	cttnccactn	ctcattcggg	tnatgtgtct	caattaaatc	360
ccacngatgg	gggtcatttt	tntcnnttag	ccagtttgtg	nagttccgtt	attganaaaa	420
ccan						424

<210> 12

<211> 426

<212> DNA

<213> Homo sapien

<220>

<221> misc_feature

<222> (1)...(426)

<223> n = A,T,C or G

<400> 12

tttttttttt	tttttcttaa	aagctttttat	ctcctgctta	cattacccat	ctgttcttgc	60
atgttgtctg	ctttttccac	tagagccctt	aacaacttaa	tcattggttat	tttaagggct	120
ctaataattc	cnaaactggg	atcataaata	agtctcgctt	tnatgcttgt	tttctctcta	180
tcacactgtg	ttingttgctt	tttnacatgc	tttgtaattt	ttggctgaaa	gctgaaaaat	240
nacatacctg	gttntacaac	ctgaggtaan	cagccttnta	gtgtgagggt	ttatatntta	300
ctggctaaga	gctnngcnct	gttnantant	tggtgtanct	ntatatgcca	naggctttna	360
tttcnctng	tgctcttgct	tnagtacccc	attnttttag	gggttcccta	naaactctat	420
ctnaat						426

<210> 13

<211> 419

<212> DNA

<213> Homo sapien

<220>

<221> misc_feature

<222> (1)...(419)

<223> n = A,T,C or G

<400> 13

tttttttttt	tttttnagat	agactctcac	tctttcgccc	aggctggagt	gcagtggcgc	60
aatcaaggct	cactgcaacc	tctgccttat	aaagcatttn	ctaaagggtac	aagctaaatt	120
ttaaaaatat	ctctncacaa	ctaattgtata	acaaaaatta	gttctacctc	ataaacnct	180
ggctcagccc	tcgnaacaca	tttcctgtt	ctcaactgat	gaacactcca	naaacagAAC	240
anatntaagc	ttttccaggc	ccagaaaagc	tcgcgagggg	atttgctntg	tgtgtgacac	300
acttgccacc	ctgtggcagc	acagctccac	acntgctttg	ggccgcattt	gcaagttctc	360
tgtaancccc	ctgnaagacc	cggatcagct	gggtngaaat	tgcangcnct	cttttggca	419

<210> 14

<211> 400

<212> DNA

<213> Homo sapien

<220>

<221> misc_feature

<222> (1)...(400)

<223> n = A,T,C or G

<400> 14

aanccattgc	caaggggtatc	cggaggattg	tggctgtcac	aggtncgag	gcccanaagg	60
ccctcaggaa	agcaaagagc	ttgaaaaatg	tctctctgtc	atggaagccn	aagtgaaggc	120
tcanactgct	ccaacaagga	tntgcanagg	gagatcgcta	accttggaga	ggccctggcc	180
actgcagtcn	tcccccantg	gcagaaggat	gaattgcggg	agactctcan	atcccttang	240
gaaggctcgtg	gatnacttgg	accgagcctc	nnaagccaat	ntccagaaca	agtgttggag	300
aagacaaagc	anttcacga	cgccaacccc	naccggcctc	tnttctcctg	ganattgana	360
gcggcgcccc	cgcccagggc	cttaataanc	cntgaagctn			400

<210> 15

<211> 395

<212> DNA

<213> Homo sapien

<220>

<221> misc_feature

<222> (1)...(395)

<223> n = A,T,C or G

<400> 15

tgctttgctg	cgtccaggaa	gattagatng	aanaatacat	attgatttgc	caaataaaca	60
agcgagatta	gacntactga	anatccatgc	aggtcccatt	acaaagcatg	gtgaaataga	120
tgatgaagca	attgtgaagc	tatcggaagg	ctttnatgga	gcagatctga	gaaatgtttg	180
tactgaagca	ggtatgttcg	caattcgtgc	tgatcatgat	ttttagtagt	aggaagactt	240
catgaaagcn	gtcagaanag	tggctnatte	tnaaagctgg	agtctaaatt	ggacnacnac	300
ctntgtatatt	actgttggan	ttttgatgct	gcatgacaga	ttttgcttan	tgtaaaaatn	360
aagttcaaga	aaattatggt	agttttggcc	attat			395

<210> 16

<211> 404

<212> DNA

<213> Homo sapien

<220>

<221> misc_feature

<222> (1)...(404)

<223> n = A,T,C or G

<400> 16

ccaccactaa	aatcctggct	gagccctacn	agtagctgtg	cccctcccc	aggacgagat	60
nagggcacac	cctttaagtn	aggtgacagg	tcacctttaa	gtgaggacag	tcagctnaat	120
ttcacctctt	gggcttgagt	acctggttct	cgtgcccctga	ggcgacnctn	agccctgcag	180
ctnccatgta	cgtgctgcc	atngtcttga	tcttctccac	gccnctnaac	ttgggcttca	240
gtaggagctg	caggcnagaa	ngaagcgggt	aacagcgcca	ctccatagcc	gcagcnggc	300
tgcccctgct	tctcaaggag	gggtgtgggg	ttcctccacc	atcgccgccc	ttgcaaacac	360
ntctcanggc	ttccctnccg	gctnancgca	ngacttaagc	atgg		404

<210> 17

<211> 360
 <212> DNA
 <213> Homo sapien

<220>
 <221> misc_feature
 <222> (1)...(360)
 <223> n = A,T,C or G

<400> 17
 ggccagaagc tttccacaaa ccagtgaagg tggcagcaaa gaaagcctct tagacnagga 60
 gctggcagca gctgctatct ngatngaeng cagaaaccaa ccactaatc agcaaacaca 120
 acctcatacc tnaccgcttc cctttnaatg gccttcggtg tgtgcgcaca tgggcacgtg 180
 cggggagaac catacttatt cccctnttcc cggcctacca cctctnctcc cccttctctt 240
 ctctncaatt actntctccn ctgctttntt ctanaccta ctgctngtnt cnanagccng 300
 cccgcaatta cctggcaaaa ctgcgcaccc ttcgggcagc gctaaanaat gcacatttac 360

<210> 18
 <211> 316
 <212> DNA
 <213> Homo sapien

<220>
 <221> misc_feature
 <222> (1)...(316)
 <223> n = A,T,C or G

<400> 18
 atacatatac acatatatga ttttagatag agccatatac ctngaagtag tanatttggt 60
 tgttgtgtata tgtatgtgtc tactcatttt aaataaactt gtgatagaga tgtaattntg 120
 agccagtttt tcatttgctt aaatnactca ccaagtaact aattaagtn tctttactct 180
 taatgttnag tagtgagatt ctgttgaagg tgatattaaa aaccattcta tattaattaa 240
 cattcatgtt gttttttaaa agcttatttg aaatcnaatt atgattattt ttcataccag 300
 tcgatnttat gtangt 316

<210> 19
 <211> 350
 <212> DNA
 <213> Homo sapien

<220>
 <221> misc_feature
 <222> (1)...(350)
 <223> n = A,T,C or G

<400> 19
 aagggatgca nataatgctg tgtatgagct tgatggaaaa gaactctgta gtgaaaggggt 60
 tactattgaa catgctnggg ctgcggcacg aggtggaaga ggtagaggac gatactctga 120
 ccgttttagt agtcgcagac ctcgaaatga tagacgaaat gctccacctg taagaacaga 180
 anacgtctt atagttgaga atttatcttc aagagtcagc tggcagggtt gttganatac 240
 agttttgagt tnttttgatg tggcttttta aaaaagttat gggttactna tggtatattg 300
 ttttattaaa agtagttttn aattaatgga tntgatggaa ttgttggttt 350

<210> 20
 <211> 367

<212> DNA
 <213> Homo sapien

<220>
 <221> misc_feature
 <222> (1)...(367)
 <223> n = A,T,C or G

<400> 20
 gntnnnncna agatcctnct ntccccngg gngcccccnc cncngtnat naccggtttn 60
 ntaanatcnn gccgcncccg aagtctcnc nntgccgaga tgnccttat ncnncnatgn 120
 ncaattntga cctnnggcga anaatggcng nngtgtatca gtntccnctc tngngnctct 180
 tagnatctga ccaactangac ccnctatcct ctcaaaccct gtanncngcc ctaatttggtg 240
 ccaatttagtg catgntanag cntcctggcc cagatggcnt ccatatcctg gtncggcttc 300
 cgcccctacc angncatccn catctactag agcttatccg ctncntgngg cgcaccggnt 360
 ccccnct 367

<210> 21
 <211> 366
 <212> DNA
 <213> Homo sapien

<220>
 <221> misc_feature
 <222> (1)...(366)
 <223> n = A,T,C or G

<400> 21
 cccaacacaa tgggtctaagt anaactgtat tgctctgtag tatagttcca cattggcaac 60
 ctacaatggg aaaatccata cataagtcag ttacttcctn atgagctttc tccttctgaa 120
 tcctttatct tctgaagaaa gtacacacct tggtnatgat atctttgaat tgcccttctt 180
 tccaggcatc agttggatga ttcacatggg taattatggc attatcatat tcttcatact 240
 tgtcatacga aaacaccagt tctgcccna gatgagcttg ttctgcagct ctttagcacct 300
 tgggaatatt cactctagac cagaaacagc tcccggtgct ccctcatttt ctgaggctta 360
 aatttn 366

<210> 22
 <211> 315
 <212> DNA
 <213> Homo sapien

<220>
 <221> misc_feature
 <222> (1)...(315)
 <223> n = A,T,C or G

<400> 22
 acttaatgca atctctggag gataatttgg atcaagaaat aaagaanaaa tgaattagga 60
 gaagaaatna ctgggtnata tttcaatatt ttagaacttt aanaatgttg actatgattt 120
 caatatatct gtnaaaactg agatacanag ttgacctata tctgcatttt gataattaaa 180
 cnaatnnatt ctatttnaat gttgtttcag agtcacagca cagactgaaa ctttttttga 240
 atacctnaat atcacacttn tncttnnaat gatgttgaag acaatgatga catgccttna 300
 gcatataatg tcgac 315

<210> 23

<211> 202
 <212> DNA
 <213> Homo sapien

<220>
 <221> misc_feature
 <222> (1)...(202)
 <223> n = A,T,C or G

<400> 23
 actaatccag tgtggtgnaa ttccattgtg ttgggcaact caggatatta aatttatnat 60
 ttaaaaattc ccaagagaaa naaactccag gccctgattg tttcactggg gaattttacc 120
 aaatgttnca nnaaganatg acgctgattc tgtnaaatct ttttcagaag atagaggaga 180
 acaccaccg ntccatttta tg 202

<210> 24
 <211> 365
 <212> DNA
 <213> Homo sapien

<220>
 <221> misc_feature
 <222> (1)...(365)
 <223> n = A,T,C or G

<400> 24
 ggattttcttg cctttttctc cctttttaag tatcaatgta tgaaatccac ctgtaccacc 60
 ctttctgcca tacaaccgct accacatctg gctcctagaa cctgttttgc tttcatagat 120
 ggatctcgga accnagtgtt nacttcattt ttaaacccca ttttagcaga tngtttgctn 180
 tggctctgtc gtattcacca tggggcctgt acacaccacg tgtggttata gtcaaacaca 240
 gtgccctcca ttgtggccac atgggagacc catnaccna tactgcatcc tgggctgatn 300
 acggcactgc atctnaccg acntgggatt gaaccggggg tgggcagcng aattgaacag 360
 gatca 365

<210> 25
 <211> 359
 <212> DNA
 <213> Homo sapien

<220>
 <221> misc_feature
 <222> (1)...(359)
 <223> n = A,T,C or G

<400> 25
 gtttcttgct tcaacagtgc ttggacggaa cccggcgctc gttccccacc ccggccggcc 60
 gcccatagcc agccctccgt cacctcttca ccgcaccctc ggactgcccc aaggcccccg 120
 ccgcnctcc ngcgcnccgc agccaccgcc gccnccncca cctctccttn gtcccgcct 180
 nacaacgcgt ccacctcgca ngttcgccng aactaccacc nggactcata ngccgcctc 240
 aaccgcccga tcaacctgga gctctncccc ccgacnttaa cctttccntg tcttacttac 300
 nttaaccgcc gnttatcttg cttnaaaaga acttttcccc aatactttct ttcacnnt 359

<210> 26
 <211> 400
 <212> DNA

```

<213> Homo sapien

<220>
<221> misc_feature
<222> (1)...(400)
<223> n = A,T,C or G

<400> 26
agtgaacag tatatgtgaa aaggagtttg tgannagcta cataaaaata ttagatatct      60
ttataatttc caataggata ctcatcagtt ttgaataana gacatattct agagaaacca      120
ggtttctggt ttcagatttg aactctcaag agcttggag ttatcactcc catcctcacg      180
acnacnaana aatctnaacn aacngaana caatgacttt tcttagatct gtcaaagaac      240
ttcagccacg aggaaaacta tcncctnaa tactggggac tggaaagaga gggtagacag      300
aatcacagt aatcatagcc caagatcagc ttgcccgag ctnaagctng tacgatnatt      360
acttacagg accacttcac agtnngtnga tnaantgccn      400

<210> 27
<211> 366
<212> DNA
<213> Homo sapien

<220>
<221> misc_feature
<222> (1)...(366)
<223> n = A,T,C or G

<400> 27
gaatttctta gaaactgaag tttactctgt tccaagatat atcttcactg tcttaatcaa      60
agggcgctng aatcatagca aatattctca tctttcaact aactttaagt agtntcctg      120
gaattttaca tttccagaa aacactcctt tctgtatctg tgaaagaaag tgtgcctcag      180
gctgtagact gggctgcact ggacacctgc gggggactct ggctnagtgn ggacatggtc      240
agtattgatt ttccctcanac tcagcctgtg tagctntgaa agcatggaac agattacact      300
gcagttnacg tcatcccaca catcttgga cccnagaccc ggggaggtca catagtccgt      360
tatgna      366

<210> 28
<211> 402
<212> DNA
<213> Homo sapien

<220>
<221> misc_feature
<222> (1)...(402)
<223> n = A,T,C or G

<400> 28
agtgggagcc tcctccttcc ccactcagtt ctttacatcc ccgaggcgca gctgggcnaa      60
ggaagtggcc agctgcagcg cctcctgcag gcagccaacg ttcttgcttg tggcctgtgc      120
agacacatcc ttgccaccac ctttaccgtc catcangcct gacacctgct gcaccactc      180
gctngctttt aagccccgat nggctgcatt ctgggggact tgacacaggc ncgtgatctt      240
gccagcctca ttgtccaccg tgaagagcat ggcaaaaagt ctgaggggag tgcattctga      300
anagcttcaa ggcttcattc agggccttng ctnaggcgcc nctctccatc tccnggaata      360
acnagaggct ggttnnggtn actntcaata aactgcttcg tc      402

<210> 29

```

<211> 175
 <212> DNA
 <213> Homo sapien

<400> 29
 cggacgggca tgaccgggtcc ggtcagctgg gtggccagtt tcagttcttc agcagaactg 60
 tctcccttct tggggggccga gggcttcctg ggaagagga tgagtttggg gcggtactcc 120
 ttcagccgct gcacgttggt ctgcaggac tccgtggact tgtccgcct cctcg 175

<210> 30
 <211> 360
 <212> DNA
 <213> Homo sapien

<400> 30
 ttgtatttct tatgatctct gatgggttct tctcgaaaat gccagtgga agactttgtg 60
 gcatgctcca gatttaaadc cagctgaggc tccctttgtt ttcagttcca tgtaacaadc 120
 tggaaggaaa cttcacggac aggaagactg ctggagaaga gaagcgtgtt agcccatattg 180
 aggtctgggg aatcatgtaa agggtagcca gacctactt ttagttattt acatcaatga 240
 gttctttcag ggaaccaaac ccagaattcg gtgcaaaagc caaacatctt ggtgggattt 300
 gataaatgcc ttgggacctg gagtgcctggg cttgtgcaca ggaagagcac cagccgctga 360

<210> 31
 <211> 380
 <212> DNA
 <213> Homo sapien

<220>
 <221> misc_feature
 <222> (1) ... (380)
 <223> n = A,T,C or G

<400> 31
 acgctctaag cctgtccacg agctcaatag ggaagcctgt gatgactaca gactttgcca 60
 acgctacgcc atggtttatg gatacaatgc tgcctataan cgctacttca ggaagcgccg 120
 agggaccnaa tgagactgag ggaagaaaaa aaatctcttt tttctggag gctggcacct 180
 gattttgtat cccctgtttn cagcattncn gaaatacata ggcttatata caatgcttct 240
 ttctgtata ttctctgtc ttgctgcacc ccttnttccc gccccagat tgataagtaa 300
 tgaaagtga ctgcagtnag ggtcaangga gactcancat atgtgattgt tccntnataa 360
 acttctggtg tgatactttc 380

<210> 32
 <211> 440
 <212> DNA
 <213> Homo sapien

<220>
 <221> misc_feature
 <222> (1) ... (440)
 <223> n = A,T,C or G

<400> 32
 gtgtatggga gccctgact cctcacgtgc ctgatctgtg cccttgggtc caggtcaggc 60
 ccacccctg cacctccacc tgccccagcc cctgcctctg cccaagtgg ggccagctgc 120
 cctcacttct ggggtggatg atgtgacctt cctnggggga ctgcggaagg gacaagggtt 180

```
ccctgaagtc ttacgggtcca acatcaggac caagtcccat ggacatgctg acaggggtccc 240
caggggagac cgtntcanta gggatgtgtg cctggctgtg tacgtgggtg tgcagtgcac 300
gtganaagca cgtggcggct tctggggggc atgtttggg aaggaagtgt gcccncacc 360
cttggagaac ctacgtcccn gtagccccc gccctggcac agcngcatnc acttcaaggg 420
cacccttttg gggttgggtg 440
```

```
<210> 33
<211> 345
<212> DNA
<213> Homo sapien
```

```
<220>
<221> misc_feature
<222> (1)...(345)
<223> n = A,T,C or G
```

```
<400> 33
tattttaaca atgtttatta ttcatttatc cctctataga accaccaccc acaccgagga 60
gattatttgg agtgggtccc aacctagggc ctggactctg aaatctaact cccacttcc 120
ctcattttgt gacttaggtg ggggcatggg tcagtcagaa ctggtgtctc ctattggatc 180
gtgcagaagg aggacctagg cacacacata tgggtggccac acccaggagg gttgattggc 240
aggctggaag acaaaagtct cccaataaag gcacttttac ctcaaagang ggggtgggagt 300
tggtctgctg ggaatgttgt tgttgggggtg gggaagantt atttc 345
```

```
<210> 34
<211> 440
<212> DNA
<213> Homo sapien
```

```
<220>
<221> misc_feature
<222> (1)...(440)
<223> n = A,T,C or G
```

```
<400> 34
tgtaattttt ttattggaaa acaaataaac aacttggaaat ggattttgag gcaaattgtg 60
ccataagcag attttaagtg gctaaacaaa gtttaaaaag caagtaacaa taaaagaaaa 120
tgtttctggg acaggaccag cagtacaaaa aaatagtgtg cgagtacctg gataatacac 180
ccgttttgca atagtgcac ttttaagtac atattgttga ctgtccatag tccacgcaga 240
gttacaactc cacacttcaa caacaacatg ctgacagttc ctaaagaaaa ctactttaaa 300
aaaggcataa cccagatgtt ccttcatttg accaactcca tctnagttta gatgtgcaga 360
agggtttana ttttcccaga gtaagccnca tgcaacatgt tacttgatca attttctaaa 420
ataagggttt aggacaatga 440
```

```
<210> 35
<211> 540
<212> DNA
<213> Homo sapien
```

```
<220>
<221> misc_feature
<222> (1)...(540)
<223> n = A,T,C or G
```

```
<400> 35
```

atagatggaa	tttattaagc	ttttcacatg	tgatagcaca	tagttttaat	tgcacccaaa	60
gtactaacia	aaactctagc	aatcaagaat	ggcagcatgt	tattttataa	caatcaaacac	120
ctgtggcttt	taaaatttgg	ttttcataag	ataatttata	ctgaagtaaa	tctagccatg	180
cttttaaaaa	atgcttttagg	tcactccaag	cttggcagtt	aacatttggc	ataaacaata	240
ataaaacaat	cacaatttta	taaataacaa	atacaacatt	gtaggccata	atcatatata	300
gtataagggg	aaagggtgga	gtgttganta	agcagttatt	agaatagaat	accttggcct	360
ctatgcaaat	atgtctagac	actttgattc	actcagccct	gacattcagt	tttcaaagtt	420
aggaaacagg	ttctacagta	tcattttaca	gtttccaaca	cattgaaaac	aagtagaaaa	480
tgatganttg	atttttatta	atgcattaca	tcctcaagan	ttatcaccaa	cccctcaggt	540

<210> 36
 <211> 555
 <212> DNA
 <213> Homo sapien

<220>
 <221> misc_feature
 <222> (1)...(555)
 <223> n = A,T,C or G

cttcgtgtgc	ttgaaaattg	gagcctgccc	ctcggcccat	aagcccttgt	tgggaactga	60
gaagtgtata	tggggcccaa	nctactggtg	ccagaacaca	gagacagcag	cccantgcaa	120
tgctgtcgag	cattgcaaac	gccatgtgtg	gaactaggag	gaggaatatt	ccatcttggc	180
agaaaccaca	gcattgggtt	ttttctactt	gtgtgtctgg	gggaatgaac	gcacagatct	240
gtttgacttt	gttataaaaa	tagggctccc	ccacctcccc	cntttctgtg	tnctttattg	300
tagcantgct	gtctgcaagg	gagcccctan	cccctggcag	acananctgc	ttcagtgcc	360
ctttcctctc	tgctaaatgg	atgttgatgc	actggaggct	ttttancttg	cccttgcatg	420
gcncctgctg	gaggaagana	aaactctgct	ggcatgaccc	acagtttctt	gactggangc	480
cntcaaccct	cttggttgaa	gccttggtct	gacctgaca	tnctgttggg	cnctgggtng	540
gnctgggctt	ctnaa					555

<210> 37
 <211> 280
 <212> DNA
 <213> Homo sapien

<220>
 <221> misc_feature
 <222> (1)...(280)
 <223> n = A,T,C or G

ccaccgacta	taagaactat	gccctcgtgt	attcctgtac	ctgcatcatc	caactttttc	60
acgtggattt	tgcttggtac	ttggcaagaa	accctaattc	ccctccagaa	acagtggact	120
ctctaaaaaa	tatcctgact	tctaataaca	ttgatntcaa	gaaaatgacg	gtcacagacc	180
agggtgaactg	ccccnagctc	tcgtaaccag	gttctacagg	gaggctgcac	ccactccatg	240
ttntctctgc	ttcgctttcc	cctaccccac	cccccgccat			280

<210> 38
 <211> 303
 <212> DNA
 <213> Homo sapien

<220>

<221> misc_feature
<222> (1)...(303)
<223> n = A,T,C or G

<400> 38
catcgagctg gttgtcttct tgcctgccct gtgtcgtaaa atgggggtcc cttactgcat 60
tatcaagggg aaggcaagac tgggacgtct agtccacagg aagacctgca ccaactgtcgc 120
cttcacacag gtgaactcgg aagacaaagg cgctttggct nagctggtnn aagctatcag 180
gaccaattac aatgacngat acgatnagat ccgccntcac tggggtagca atgtcctggg 240
tcctaagtct gtggctcgta tcgccnagct cgaanaggcn aangctaaag aacttgccac 300
taa 303

<210> 39
<211> 300
<212> DNA
<213> Homo sapien

<220>
<221> misc_feature
<222> (1)...(300)
<223> n = A,T,C or G

<400> 39
gactcagcgg ctgggtgctct tctgtgacac aagcccagca ctccagggtcc caaggcattt 60
atcaaattccc accaagatnt ttggcttttg caccgaattc tgggttttgt tccctnaaag 120
aactcattga tgtaatnac tnaaagttag gtctgggtac cctttacatg attccccaga 180
cctcanatgg gctaacacgc ttctcttctc cagcagttct cctntccgtg aagttaacct 240
ccagattggt acatggaact gaanacaaag ggagccctcag ctngatttaa atctggagca 300

<210> 40
<211> 318
<212> DNA
<213> Homo sapien

<220>
<221> misc_feature
<222> (1)...(318)
<223> n = A,T,C or G

<400> 40
cccaacacaa tgggtgagga caaatcagtt ctctgtgacc agacatgaga aggttgccaa 60
tgggctgttg ggcgaccaag gccttcccgg agtcttcgtc ctctatgagc tctcgcccat 120
gatgggtgaag ctgacggaga agcacagggtc cttcaccac ttcttgacag gtgtgtgcgc 180
catcattggg ggcattgtca cagtgggtgg actcatcgat tcgctcatct accactcagc 240
acgagccatc cagaaaaaaa ttgatctngg gaagacnacg tagtcaccct cggtntcttc 300
tctgtctcct ctttctcc 318

<210> 41
<211> 302
<212> DNA
<213> Homo sapien

<220>
<221> misc_feature
<222> (1)...(302)

```

<223> n = A,T,C or G

<400> 41
acttagatgg ggtccgttca ggggatacca gcgttcacat ttttcctttt aagaaaggg      60
cttggcctga atgttcccca tccggacaca ggctgcatgt ctctgtnagt gtcaaagctg      120
ccatnaccat ctcggtaacc tactcttact ccacaatgtc tatnttcaact gcagggctct      180
ataatnagtc cataatgtaa atgcctggcc caagacntat ggcctgagtt tatccnaggc      240
ccaaacnatt accagacatt cctcttanat tgaaaacgga tntctttccc ttggcaaaga      300
tc                                                                    302

<210> 42
<211> 299
<212> DNA
<213> Homo sapien

<220>
<221> misc_feature
<222> (1) ... (299)
<223> n = A,T,C or G

<400> 42
cttaataagt ttaaggccaa ggcccgttcc attcttctag caactgacgt tgccagccga      60
ggtttggaca tacctcatgt aaatgtggtt gtcaactttg acattcctac ccattccaag      120
gattacatcc atcgagtagg tcgaacagct agagctgggc gtcccggaaa ggctattact      180
tttgtcacac agtatgatgt ggaactcttc cagcgcatag aacacttnat tgggaagaaa      240
ctaccaggtt ttccaacaca ggatgatgag gttatgatgc tnacggaacg cgtcgctna      299

<210> 43
<211> 305
<212> DNA
<213> Homo sapien

<220>
<221> misc_feature
<222> (1) ... (305)
<223> n = A,T,C or G

<400> 43
ccaacaatgt caagacagcc gtctgtgaca tcccacctcg tggcctcaan atggcagtca      60
ccttcattgg caatagcaca gccntccggg agctcttcaa gcgcctctcg gagcagttca      120
ctgccatggt ccgccggaag gccttctctc actggtacac aggcgagggc atggacaaga      180
tggagttcac cgaggctgag agcaacatga acgacctcgt ctctnagtat cagcagtacc      240
gggatgccac cgcagaaana ggaggaggat ttcggtnagg aggccgaaga aggaggcctg      300
aggca                                                                    305

<210> 44
<211> 399
<212> DNA
<213> Homo sapien

<220>
<221> misc_feature
<222> (1) ... (399)
<223> n = A,T,C or G

```

```

<400> 44
tttctgtggg ggaaacctga tctcgacnaa attagagaat tttgtcagcg gtatttcggc      60
tggaacagaa cgaaaacnga tnaatctctg tttcctgtat taaagcaact cgatncccag      120
cagacacagc tccnaattga ttccttcttt ngattagcac aacagggaga aagaanatgc      180
ttaacgtatt aagagccnga gactaaacag agctttgaca tgratgctta ggaaagagaa      240
agaagcagcn gccgcgnaa ttngaagcng tttctgttgc cntgganaaa gaatttgagc      300
ttctttatta ggccaacgaa aaaccccgaa ananaggcnt tacnatacct tngaaaantc      360
tccngccnna aaaagaaaga agctttcnga ttcttaacc      399

```

```

<210> 45
<211> 440
<212> DNA
<213> Homo sapien

```

```

<220>
<221> misc_feature
<222> (1)...(440)
<223> n = A,T,C or G

```

```

<400> 45
gcgggagcag aagctaaagc caaagcccaa gagagtggca gtgccagcac tggtgccagt      60
accagtacca ataacagtgc cagtgccagt gccagcacca gtgggtggctt cagtgcctgg      120
gccagcctga ccgccactct cacatttggg ctcttcgctg gccttgggtg agctggtgcc      180
agcaccagtg gcagctctgg tgcctgtggg ttctcctaca agtgagattt taggtatctg      240
ccttggtttc agtggggaca tctggggctt anggggcngg gataaggagc tggatgattc      300
taggaaggcc cangttggag aangatgtgn anagtgtgcc aagacactgc ttttggcatt      360
ttattccttt ctgtttgctg gangtcaatt gacccttnna ntttctctta cttgtgtttt      420
canatatngt taatcctgcc      440

```

```

<210> 46
<211> 472
<212> DNA
<213> Homo sapien

```

```

<220>
<221> misc_feature
<222> (1)...(472)
<223> n = A,T,C or G

```

```

<400> 46
gctctgtaat ttcacatttt aaaccttccc ttgacctcac attcctcttc ggccacctct      60
gtttctctgt tcctcttcac agcaaaaact gttcaaaaaga gttgttgatt actttcattt      120
ccactttctc acccccattc tcccctcaat taactctcct tcatcccat gatgccatta      180
tgtggctntt attanagtca ccaaccttat tctccaaaac anaagcaaca aggactttga      240
cttctcagca gcactcagct ctggtncttg aaacaccccc gttacttgct attcctccta      300
cctcataaca atctccttcc cagcctctac tgetgccttc tctgagttct tcccagggtc      360
ctaggctcag atgtagtgtg gctcaaccct gctacacaaa gnaatctcct gaaagcctgt      420
aaaaatgtcc atncntgtcc tgtgagtgtat ctncangna naataacaaa tt      472

```

```

<210> 47
<211> 550
<212> DNA
<213> Homo sapien

```

```

<220>

```

<221> misc_feature
<222> (1) ... (550)
<223> n = A,T,C or G

<400> 47
ccttcctccg cctggccatc cccagcatgc tcatgctgtg catggagtgg tgggcctatg 60
aggctcgggag cttcctcagt ggtctgtatg aggatggatg acggggactg gtgggaacct 120
gggggcccctg tctgggtgca aggcgacagc tgtctttctt caccaggcat cctcggcatg 180
gtggagctgg gcgctcagtc catcgtgtat gaactggcca tcattgtgta catggtccct 240
gcaggcttca gtgtggctgc cagtgtccgg gtangaaacg ctctgggtgc tggagacatg 300
gaagcaggca cggaagtccct ctaccgtttc cctgctgatt acagtgtctt ttgctgtanc 360
cttcagtgtc ctgctgttaa gctgtaagga tcacntgggg tacattttta ctaccgaccg 420
agaacatcat taatctgggtg gctcagggtg ttccaattta tgctgtttcc cacctctttg 480
aagctcttgc tgctcaggta cacgccaatt ttgaaaagta aacaacgtgc ctcggagtgg 540
gaattctgct 550

<210> 48
<211> 214
<212> DNA
<213> Homo sapien

<220>
<221> misc_feature
<222> (1) ... (214)
<223> n = A,T,C or G

<400> 48
agaaggacat aaacaagctg aacctgccca agacgtgtga tatcagcttc tcagatccag 60
acaacctcct caacttcaag ctggatcatct gtcctgatna gggcttctac nagagtggga 120
agtttgtggt cagttttaag gtgggccagg gttaccgcga tgatccccc aaggtgaagt 180
gtgagacnat ggtctatcac cccnacattg acct 214

<210> 49
<211> 267
<212> DNA
<213> Homo sapien

<220>
<221> misc_feature
<222> (1) ... (267)
<223> n = A,T,C or G

<400> 49
atctgcctaa aatttattca aataatgaaa atnaatctgt ttttaagaaat tcagtctttt 60
agtttttagg acaactatgc acaaagtac gatggagaat tctttttgga tnaactctag 120
gtngaggaaac ttaatccaac cgagactntt gtgaaggta gaanacagga gagggaatct 180
tggcaaggaa tggagacnga gtttgcaaat tgcagctaga gtnaatngtt ntaaatggga 240
ctgctnttgt gtctcccang gaaagtt 267

<210> 50
<211> 300
<212> DNA
<213> Homo sapien

<220>

<221> misc_feature
 <222> (1)...(300)
 <223> n = A,T,C or G

<400> 50
 gactgggtca aagctgcatg aaaccaggcc ctggcagcaa cctgggaatg gctggaggtg 60
 ggagagaacc tgacttctct tccctctccc ctccccaac attactggaa ctctgtcctg 120
 ttgggatctt ctgagcttgt tccctgctg ggtgggacag aggacaaagg agaagggagg 180
 gtctagaaga ggcagccctt ctttgtcctc tggggtnaat gagcttgacc tanagtagat 240
 ggagagacca anagcctctg atttttaatt tccataanat gttcnaagta tatntntacc 300

<210> 51
 <211> 300
 <212> DNA
 <213> Homo sapien

<220>
 <221> misc_feature
 <222> (1)...(300)
 <223> n = A,T,C or G

<400> 51
 gggtaaaatc ctgcagcacc cactctggaa aatactgctc ttaattttcc tgaaggtggc 60
 cccctatttc tagttgtgcc aggattaggg atgtggggta tagggcattt aaatcctctc 120
 aagcgctctc caagcacccc cggcctgggg gtnagtttct catcccgtta ctgctgctgg 180
 gatcaggtnn aataaatgga actcttctct tctggcctcc aaagcagcct aaaaactgag 240
 gggctctgtt agaggggacc tccaccctnn ggaagtcga ggggctnggg aagggtttct 300

<210> 52
 <211> 267
 <212> DNA
 <213> Homo sapien

<220>
 <221> misc_feature
 <222> (1)...(267)
 <223> n = A,T,C or G

<400> 52
 aaaatcaact tcntgcatta atanacanat tctanancag gaagtgaana taattttctg 60
 cacctatcaa ggaacnnact tgattgcctc tattnaacan atatatcgag ttncataact 120
 tacctgaata ccnccgcata actctcaacc nanatnctc nccatgacac tcnttcttna 180
 atgctantcc cgaattcttc attatatcng tgatgttcgn cctgtnata tatcagcaag 240
 gtatgtncn taactgccga nncaang 267

<210> 53
 <211> 401
 <212> DNA
 <213> Homo sapien

<400> 53
 agsccttagc atcatgtaga agcaaactgc acctatggct gagataggtg caatgaccta 60
 caagattttg tgttttctag ctgtccagga aaagccatct tcagtcttgc tgacagtcaa 120
 agagcaagtg aaaccatttc cagcctaaac tacataaaag cagccgaacc aatgattaaa 180
 gacctctaag gctccataat catcattaaa tatgcccaaa ctcatgtga ctttttattt 240

tatatacagg	attaaaatca	acattaaatc	atcttattta	catggccatc	ggtgctgaaa	300
ttgagcattt	taaatagtag	agtaggctgg	tatacattag	gaaatggact	gcactggagg	360
caaatagaaa	actaaagaaa	ttagataggc	tggaaatgct	t		401

<210> 54
 <211> 401
 <212> DNA
 <213> Homo sapien

<400> 54						
cccaacacaa	tggataaaaa	cacttatagt	aaatggggac	attcactata	atgatctaag	60
aagctacaga	ttgtcatagt	tgttttcctg	ctttacaaaa	ttgctccaga	tctggaatgc	120
cagtttgacc	tttgcttctt	ataatatttc	ctttttttcc	cctctttgaa	tctctgtata	180
tttgattctt	aactaaaatt	gttctcttaa	atattctgaa	tcctggtaat	taaaagtttg	240
ggtgtatttt	ctttacctcc	aaggaaagaa	ctactagcta	caaaaaatat	tttggataaa	300
gcattgtttt	ggtataaggt	acataatttt	gttgaagaca	ccagactgaa	gtaaacagct	360
gtgcacccaa	tttattatag	ttttgtaagt	aacaatatgt	a		401

<210> 55
 <211> 933
 <212> DNA
 <213> Homo sapien

<400> 55						
tttactgctt	ggcaaagtac	cctgagcatc	agcagagatg	ccgagatgaa	atcagggaaac	60
tcctagggga	tgggtcttct	attacctggg	aacacctgag	ccagatgcct	tacaccacga	120
tgtgcatcaa	ggaatgcctc	cgctcttacg	caccggtagt	aaactatccc	ggttactcga	180
caaaccctac	acctttccag	atggacgctc	cttacctgca	ggaataactg	tgtttatcaa	240
tatttgggct	cttcaccaca	acccctattt	ctgggaagac	cctcaggtct	ttaaccctt	300
gagattctcc	agggaaaatt	ctgaaaaaat	acatccctat	gccttcatac	cattctcagc	360
tggattaagg	aactgcattg	ggcagcattt	tgccataatt	gagtgtaaag	tggcagtggc	420
attaactctg	ctccgcttca	agctggctcc	agaccactca	aggccacca	gctgtcgtca	480
agttgcctca	agtccaagaa	tggaatccat	gtgtttgcaa	aaaaagtttg	ctaattttaa	540
gtccttttct	tataagaatt	aakgagacaa	tttccttacc	aaaggaagaa	caaaaggata	600
aatataatac	aaaatatatg	tatatgggtg	tttgacaaat	tataataact	aggatacttc	660
tgactgggtt	tgacatccat	taacagtaat	tttaatttct	ttgctgtatc	tggtgaaacc	720
cacaaaaaca	cctgaaaaaa	ctcaagctga	gttccaatgc	gaagggaaat	gattgggttg	780
ggtaactagt	ggtagagtgg	ctttcaagca	tagtttgatc	aaaactccac	tcagtatctg	840
cattactttt	atctctgcaa	atctctgcat	gatagcttta	ttctcagtta	tctttcccca	900
taataaaaaa	tatctgccaa	aaaaaaaaaa	aaa			933

<210> 56
 <211> 480
 <212> DNA
 <213> Homo sapien

<400> 56						
ggctttgaag	catttttgtc	tgtgtctcct	gatcttcagg	tcaccaccat	gaagttctta	60
gcagtccctg	tactcttggg	agtttccatc	tttctggctt	ctgcccagaa	tccgacaaca	120
gctgtccag	ctgacacgta	tccagctact	ggtcctgctg	atgatgaagc	ccctgatgct	180
gaaaccatca	ctgctgcaac	cactgcgacc	actgctgctc	ctaccactgc	aaccaccgct	240
gcttctacca	ctgctcgtaa	agacattcca	gttttaccga	aatgggttgg	ggatctcccg	300
aatggtagag	tgtgtccctg	agatggaatc	agcttgagtc	ttctgcaatt	ggtcacaact	360
attcatgctt	cctgtgattt	catccaacta	cttaccttgc	ctacgatatc	ccctttatct	420
ctaatacagt	tattttcttt	caataaaaaa	ataactatga	gcaacaaaaa	aaaaaaaaaa	480

<210> 57
 <211> 798
 <212> DNA
 <213> Homo sapien

<400> 57
 agcctacctg gaaagccaac cagtcctcat aatggacaag atccaccagc tcctcctgtg 60
 gactaacttt gtgatatggg aagtgaatat agttaacacc ttgcacgacc aaacgaacga 120
 agatgaccag agtactctta accccttaga actgtttttc cttttgtatc tgcaatatgg 180
 gatggatttg ttttcatgag cttctagaaa tttcacttgc aagtttattt ttgcttcctg 240
 tgttactgcc attcctattt acagtatatt tgagtgaatg attatatttt taaaaagtta 300
 catggggcctt ttttggttgt cctaaactta caaacattcc actcattctg tttgtaactg 360
 tgattataat ttttgtgata atttctggcc tgattgaagg aaatttgaga ggtctgcatt 420
 tatatatttt aaatagattt gataggtttt taaattgctt tttttcataa ggtattttata 480
 aagttatttg gggttgtctg ggattgtgtg aaagaaaatt agaaccctgc tgtattttaca 540
 tttaccttgg tagtttattt gtggatggca gttttctgta gttttgggga ctgtggtagc 600
 tcttgatttg ttttgcataa tacagctgaa atctgtgtca tggattaaac tggcttatgt 660
 ggctagaata ggaagagaga aaaaatgaaa tgggtgttta ctaattttat actcccatta 720
 aaaattttta atgttaagaa aaccttaaat aaacatgatt gatcaatatg gaaaaaaaaa 780
 aaaaaaaaaa aaaaaaaaaa 798

<210> 58
 <211> 280
 <212> DNA
 <213> Homo sapien

<400> 58
 ggggcagctc ctgaccctcc acagccacct ggtcagccac cagctggggc aacgaggggtg 60
 gaggtccccc tgagcctctc gcctgcccc gccactcgtc tgggtgctgt tgatccaagt 120
 cccctgcctg gtccccaca aggactccca tccaggcccc ctctgccttg ccccttgctca 180
 tggaccatgg tctgtaggaa gggctcatgc cccttattta tgggaaccat ttcatttctaa 240
 cagaataaac cgagaaggaa accagaaaaa aaaaaaaaaa 280

<210> 59
 <211> 382
 <212> DNA
 <213> Homo sapien

<400> 59
 aggcgggagc agaagctaaa gccaaagccc aagagagtgg cagtgccagc actgggtgcca 60
 gtaccagtac caataacagt gccagtcca gtgccagcac cagtgggtggc ttcagtgtctg 120
 gtgccagcct gaccgccact ctcacatttg ggctcttcgc tggccttggg ggagctgggtg 180
 ccagcaccag tggcagctct ggtgcctgtg gtttctccta caagtgagat tttagatatt 240
 gttaatectg ccagtctttc tcttcaagcc aggggtgcac ctcagaaacc tactcaacac 300
 agcactctag gcagccacta tcaatcaatt gaagttgaca ctctgcatta aatctatttg 360
 ccattaaaaa aaaaaaaaaa aa 382

<210> 60
 <211> 602
 <212> DNA
 <213> Homo sapien

<400> 60
 tgaagagccg cgcggtggag ctgctgcccc atgggactgc caaccttgcc aagctgcagc 60

```

ttgtggtgga gaatagtgcc cagcgggtca tccacttggc gggtcagtgg gagaagcacc 120
gggtcccatc ctcgtgagta ccgccactcc gaaagctgca ggattgcaga gagctggaat 180
cttctcgacg gctggcagag atccaagaac tgcaccagag tgtccgggcg gctgctgaag 240
aggcccgag gaaggaggag gtctataaagc agctgatgtc agagctggag actctgccc 300
gagatgtgtc ccggctggcc tacaccagc gcatcctgga gatcgtgggc aacatccgga 360
agcagaagga agagatcacc aagatcttgt ctgatacgaa ggagcttcag aaggaaatca 420
actccctatc tgggaagctg gaccggacgt ttgcggtgac tgatgagctt gtgttcaagg 480
atgccaagaa ggacgatgct gttcgggaagg cctataagta tctagctgct ctgcacgaga 540
actgcagcca gctcatccag accatcgagg acacaggcac catcatgcgg gaggttcgag 600
ac 602

```

```

<210> 61
<211> 1368
<212> DNA
<213> Homo sapien

<220>
<221> misc_feature
<222> (1)...(1368)
<223> n = A,T,C or G

```

```

<400> 61
ccagtgcgagc cgcgtaatac gactcactat agggcgaatt gggtagcggg cccccctcg 60
agcggccgccc cttttttttt tttttttatt gatcagaatt caggctttat tattgagcaa 120
tgaaaacagc taaaacttaa ttccaagcat gtgtagttaa agtttgcaaa gtgggatatt 180
gttcacaaaa cacattcaat gtttaaacac tatttatttg aagaacaaaa tatattttaa 240
attgtttgct tctaaaaagc ccatttccct ccaagtctaa actttgtaat ttgatattaa 300
gcaatgaagt tattttgtac aatctagtta aacaagcaga atagcactag gcagaataaa 360
aaattgcaca gacgtatgca attttccaag atagcattct ttaaattcag ttttcagctt 420
ccaaagattg gttgcccata atagacttaa acatataatg atggctaaaa aaaataagta 480
tacgaaaatg taaaaaagga aatgtaagtc cactctcaat ctcataaaag gtgagagtaa 540
ggatgctaaa gcaaaataaa tgtagggttct ttttttctgt ttccgtttat catgcaatct 600
gcttctttga tatgccttag ggttacccat ttaagttaga ggttgtaatg caatgggtggg 660
aatgaaaatt gatcaaatat acaccttgtc atttcatttc aaattgcggg ctggaaactt 720
ccaaaaaag ggtaggcatg aagaaaaaaa aaatcmaatc agaacctctt cagggggttg 780
kgktctgata tggcagacar gatacaagtc ccaccaggag atggagcaat tcaaaataag 840
gtaaatgggc tgacaaggta ttattgccag catgggacag aatgagcaac aggctgaaaa 900
gtttttggat tatatagcac cttaggtctc tgatgtaggg aatttttgtt agtcaaacat 960
acgctaaact tccaaggga aatctttcag gtagecctaag cttgcttttc tagagtgatg 1020
agttgcattg ctactgtgat tttttgaaaa caaactgggt ttgtacaagt gagaaagact 1080
agagagaaaag attttagtct gtttagcaga agccatttta tctgcgtgca catggatcaa 1140
tatttctgat cccctatacc ccaggaaggg caaaatccca aagaaatgtg ttagcaaaat 1200
tggctgatgc tatcatattg ctatggacat tgatcttgcc caacacaatg gaattccacc 1260
acactggact agtggatcca ctagttctag agcggccggc caccgcggtg gagctccagc 1320
ttttgttccc tttagtgagg gtttaattgcg cgcttggcgt aatcatnn 1368

```

```

<210> 62
<211> 924
<212> DNA
<213> Homo sapien

<220>
<221> misc_feature
<222> (1)...(924)
<223> n = A,T,C or G

```


<400> 62

caaaggnaca	ggaacagctt	gnaaagtact	gncatncctn	cctgcagggg	ccagcccttt	60
gcctccaaaa	gcaataggaa	atntaaaaga	tttnactga	gaaggggncc	acgtttnart	120
tntnaatgtn	tcargnanar	tnccttncaa	atgnrcrctn	cactnactnr	gnatttggtg	180
tnccgnrtnc	mgnactatnt	caggtttgaa	aaactggatc	tgccacttat	cagttatgtg	240
accttaaaga	actccgttaa	tttctcagag	cctcagtttc	cttgtctata	agttgggagt	300
aatattaata	ctatcatttt	tccaaggatt	gatgtgaaca	ttaatgaggt	gaaatgacag	360
atgtgtatca	tggttcctaa	taaacatcca	aaatatagta	cttactattg	tcattattat	420
tacttgtttg	aagctaaaga	cctcacataa	gaatcccatc	cagcccacca	gacagagytg	480
tgagttttct	agtttggaag	agctattaaa	taacaacktc	tagtgtcaat	tctatacttg	540
ttatggtcaa	gtaactgggc	tcagcatttt	acattcattg	tctctttaag	ttctagcaat	600
gtgaagcagg	aactatgatt	atattgacta	cataaatgaa	gaaattgagg	ctcagatata	660
ttaagtaatt	ctcccagggt	cacacagcta	gaactggcaa	agcctgggat	tgatccatga	720
tcttccagca	ttgaagaatc	ataaatgtaa	ataactgcaa	ggccttttcc	tcagaagagc	780
tcttgggtgt	tgcaccaacc	cactagcact	tggtctctac	aggggaacat	ctgtgggcct	840
gggaatcact	gcacgtcgca	agagatgttg	cttctgatga	attattgttc	ctgtcagtg	900
tgtagaggca	aaaaaaaaaa	aaaa				924

<210> 63

<211> 1079

<212> DNA

<213> Homo sapien

<400> 63

agtcccaaga	actcaataat	ctcttatgtt	ttcttttgaa	gacttatttt	aaatattaac	60
tatttcggtg	cctgaatgga	aaaatataaa	cattagctca	gagacaatgg	ggtacctgtt	120
tggaatccag	ctggcagcta	taagcaccgt	tgaaaactct	gacaggcttt	gtgccctttt	180
tattaaatgg	cctcacatcc	tgaatgcagg	aatgtgttcg	tttaaataaa	cattaatctt	240
taatgttgaa	ttctgaaaac	acaaccataa	atcatagtgt	gtttttctgt	gacaatgatc	300
tagtacatta	tttccctccac	agcaaaccta	cctttccaga	aggtggaaat	tgtatttgca	360
acaatcaggg	caaaacccac	acttgaaaag	cattttacaa	tattatatct	aagttgcaca	420
gaagacccca	gtgatcacta	ggaaatctac	cacagtcacg	tttttcta	ccaagaaggt	480
caaacttcgg	ggaataatgt	gtccctcttc	tgctgctgct	ctgaaaaata	ttcgatcaaa	540
acgaagttaa	caagcagcag	ttattccaag	attagagtgc	atttgtgtat	cccatgtata	600
ctggcaatgt	ttaggtttgc	ccaaaaactc	ccagacatcc	acaatgttgt	tgggtaaacc	660
accacatctg	gtaacctctc	gatcccttag	atttgtatct	cctgcaaata	taactgtagc	720
tgactctgga	gcctcttgca	ttttctttta	aaccattttt	aactgattca	ttcgttccgc	780
agcatgccct	ctggtgctct	ccaaatggga	tgtcataagg	caaagctcat	ttcctgacac	840
attcacatgc	acacataaaa	ggtttctcat	cattttggta	cttggaagaa	gaataatctc	900
ttggcttttt	aatttcactc	ttgatttctt	caacattata	gctgtgaaat	atccttcttc	960
atgacctgta	ataatctcat	aattacttga	tctcttcttt	aggtagctat	aatatggggg	1020
aataacttcc	tgtagaaata	tcacatctgg	gctgtacaaa	gctaagtagg	aacacaccc	1079

<210> 64

<211> 1001

<212> DNA

<213> Homo sapien

<400> 64

gaatgtgcaa	cgatcaagtc	agggtatctg	tggtatccac	cactttgagc	atttatcgat	60
tctatatgtc	aggaacattt	caagttatct	gttctagcaa	ggaaatataa	aataacttata	120
gttaactatg	gcctatctac	agtgcacta	aaaactagat	tttattcctt	tccacctgtg	180
ggtttgtatt	catttaccac	cctcttttca	ttccctttct	cacccacaca	ctgtgccggg	240
cctcagggat	atactattct	actgtctgtc	tctgtaagga	ttatcatttt	agcttccaca	300

tatgagagaa	tgcattgcaaa	gtttttcttt	ccatgtcttg	cttatttcac	ttacataat	360
gacctccgct	tccatccatg	ttatttatat	tacccaatag	tgttcataaa	tatatataca	420
cacatatata	ccacattgca	tttgtccaat	tattcattga	cggaaactgg	ttaatgttat	480
atcggttgcta	ttgtggatag	tgctgcaata	aacacgcaag	tggggatata	atttgaagag	540
tttttttggt	gatgttcttc	caaattttta	gattggtttg	tctatgtttg	tgaaaatggc	600
gttagtattt	tcatagagat	tgcattgaat	ctgtagattg	ctttgggtta	gtatggttat	660
tttgatggta	ttattttttt	cattccatga	agatgagatg	tctttccatt	gtttgtgtcc	720
tctacatttt	ctttcatcaa	agttttgttg	tatttttgaa	gtagatgtat	ttcaccttat	780
agatcaagtg	tattccctaa	atattttatt	ttgtagctta	ttgtagatga	aattgccttc	840
ttgatttctt	tttcacttaa	ttcattatta	gtgtatggaa	atgttatgga	tttttatttg	900
ttgggtttta	atcaaaaact	gtattaaaact	tagagttttt	tgtggagttt	tttaagttttt	960
ctagatataa	gatcatgaca	tctacaaaaa	aaaaaaaaaa	a		1001

<210> 65

<211> 575

<212> DNA

<213> Homo sapien

<400> 65

acttgatata	aaaaggatat	ccataatgaa	tattttatac	tgcattccttt	acattagcca	60
ctaaatacgt	tattgcttga	tgaagacctt	tcacagaatc	ctatggattg	cagcattttca	120
cttggttact	tcatacccat	gccttaaaaga	ggggcagttt	ctcaaaagca	gaaacatgcc	180
gccagttctc	aagttttcct	cctaactcca	tttgaatgta	agggcagctg	gcccccaatg	240
tggggaggct	cgaacatttt	ctgaattccc	attttcttgt	tcgctggctaa	atgacagttt	300
ctgtcattac	ttagattccc	gatctttccc	aaaggtgttg	atttaciaaag	aggccagcta	360
atagccagaa	atcatgaccc	tgaaagagag	atgaaatttc	aagctgtgag	ccaggcagga	420
gctccagtat	ggcaaagggt	cttgagaatc	agccatttgg	tacaaaaaag	atttttaaaag	480
cttttatgtt	ataccatgga	gccatagaaa	ggctatggat	tgtttaagaa	ctatttttaa	540
gtgttcacga	cccaaaaagg	aaaaaaaaaa	aaaaa			575

<210> 66

<211> 831

<212> DNA

<213> Homo sapien

<400> 66

attgggctcc	ttctgctaaa	cagccacatt	gaaatggttt	aaaagcaagt	cagatcaggt	60
gatttgtaaa	attgtattta	tctgtacatg	tatgggcttt	taattcccac	caagaaagag	120
agaaattatc	tttttagtta	aaaccaaatt	tcacttttca	aaatatcttc	caacttattt	180
attggttgct	actcaattgc	ctatatatat	atatatatat	gtgtgtgtgt	gtgtgtgctc	240
gtgagcgac	gtgtgtgtat	gcgtgcgcac	gtgtgtgtat	gtgtattatc	agacataggt	300
ttctaacttt	tagatagaag	aggagcaaca	tctatgccaa	atactgtgca	ttctacaatg	360
gtgctaattc	cagacctaaa	tgatactcca	tttaatttaa	aaaagagttt	ttaaataatta	420
tctatgtgcc	tgtattttccc	ttttgagtgc	tgcaacaacat	gttaacatat	tagtgtaaaa	480
gcagatgaaa	caaccacgtg	ttctaaagtc	tagggattgt	gctataatcc	ctatttagtt	540
caaaattaac	cagaattctt	ccatgtgaaa	tggaccaaac	tcatattatt	gttatgtaaa	600
tacagagttt	taatgcagta	tgacatccca	caggggaaaa	gaatgtctgt	agtgggtgac	660
tgttatcaaa	tattttatag	aatacaatga	acggtgaaca	gactggtaac	ttgtttgagt	720
tcccatgaca	gatttgagac	ttgtcaatag	caaatcattt	ttgtatttaa	atttttgtac	780
tgatttgaaa	aacatcatta	aatatcttta	aaagtaaaaa	aaaaaaaaaa	a	831

<210> 67

<211> 590

<212> DNA

<213> Homo sapien

```

<400> 67
gtgctctgtg tattttttta ctgcattaga cattgaatag taatttgcgt taagatacgc      60
ttaaaggctc ttgtgacca tgtttccctt tgtagcaata aaatgttttt tacgaaaact      120
ttctccctgg attagcagtt taaatgaaac agagtccatc aatgaaatga gtatttataa      180
taaaaatttg ccttaaatgta tcagttcagc tcacaagtat ttaagatga ttgagaagac      240
ttgaattaaa gaaaaaaaaa ttctcaatca tattttttaa atataagact aaaattgttt      300
ttaaaacaca ttcaaataag aagtgaagtt gaactgacct tatttatact ctttttaagt      360
ttgttccttt tccctgtgcc tgtgtcaaat cttcaagtct tgctgaaaat acatttgata      420
caaagttttc tgtagttgtg ttagttcctt tgtcatgtct gtttttggct gaagaaccaa      480
gaagcagact ttctttttaa aagaattatt tctctttcaa atatttctat cctttttaa      540
aaattccttt ttatggctta tatacctaca tatttaaaaa aaaaaaaaaa      590

```

```

<210> 68
<211> 291
<212> DNA
<213> Homo sapien

<220>
<221> misc_feature
<222> (1) ... (291)
<223> n = A,T,C or G

```

```

<400> 68
gttccctttt ccggtcggcg tggctcttgcg agtggagtggt ccgctgtgcc cgggcctgca      60
ccatgagcgt cccggccttc atcgacatca gtgaagaaga tcaggctgct gagcttcgtg      120
cttarctgaa atctaaagga gctgagattt cagaagagaa ctcggaagggt ggacttcagt      180
ttgatttagc tcaaattatt gaagcctgtg atgtgtgtct gaaggaggat gataaagatg      240
ttgaaagtgt gatgaacagt ggggnatcct actcttgatc cggaanccna c      291

```

```

<210> 69
<211> 301
<212> DNA
<213> Homo sapien

<220>
<221> misc_feature
<222> (1) ... (301)
<223> n = A,T,C or G

```

```

<400> 69
tctatgagca tgccaaggct ctgtgggagg atgaaggagt gcgtgcctgc tacgaacgct      60
ccaacgagta ccagctgatt gactgtgccc agtacttcct ggacaagatc gacgtgatca      120
agcaggctga ctatgtgccg agcgatcagg acctgcttcg ctgccgtgtc ctgacttctg      180
gaatctttga gaccaagttc caggtggacn aagtcaactt ccacatgntt gacgtgggtg      240
gccagcgcga tgaacgcgcg aagtggatcc agtgcttcaa cgatgtgact gccatcatct      300
t

```

```

<210> 70
<211> 201
<212> DNA
<213> Homo sapien

```

```

<400> 70
gcggctcttc ctcgggcagc ggaagcggcg cggcggtcgg agaagtggcc taaaacttcg      60

```

```

gcgttgggtg aaagaaaatg gcccgaacca agcagactgc tcgtaagtcc accggtggga      120
aagccccccg caaacagctg gccacgaaag ccgccaggaa aagcgctccc tctaccggcg      180
gggtgaagaa gcctcatcgc t                                     201

```

```

<210> 71
<211> 301
<212> DNA
<213> Homo sapien

```

```

<220>
<221> misc_feature
<222> (1)...(301)
<223> n = A,T,C or G

```

```

<400> 71
gccggggtag tcgccgncgc cgccgccgct gcagccactg caggcaccgc tgccgccgcc      60
tgagtagtgg gcttaggaag gaagaggtca tctcgctcgg agcttcgctc ggaaggggtct    120
ttgttccttg cagccctccc acgggaatga caatggataa aagtgagctg gtacanaaag      180
ccaaactcgc tgagcaggct gagcgatatg atgatatggc tgcagccatg aaggcagtca      240
cagaacaggg gcatgaactc ttcaacgaag agagaaatct gctctctggt gcctacaaga      300
a                                     301

```

```

<210> 72
<211> 251
<212> DNA
<213> Homo sapien

```

```

<220>
<221> misc_feature
<222> (1)...(251)
<223> n = A,T,C or G

```

```

<400> 72
cttggggggg gttggggggg agactgtggg cctggaaata aaacttgtct cctctaccac      60
cacctgttac cctagcctgc acctgtccac atctctgcaa agttcagctt ccttccccag    120
gtctctgtgc actctgtctt ggatgctctg gggagctcat ggggtggagga gtctccacca    180
gagggaggct caggggactg gttggggccag ggatgaatat ttgagggata aaaattgtgt    240
aagagccaan g                                     251

```

```

<210> 73
<211> 895
<212> DNA
<213> Homo sapien

```

```

<400> 73
tttttttttt tttttcccag gccctctttt tatttacagt gataccaaac catccacttg      60
caaattcttt ggtctcccat cagctggaat taagtaggta ctgtgtatct ttgagatcat    120
gtatttgtct ccacttttgt ggatacaaga aagggaaggca cgaacagctg aaaaagaagg      180
gtatcacacc gctccagctg gaatccagca ggaacctctg agcatgccac agctgaacac      240
ttaaagaggg aaagaaggac agctgctctt catttatattt gaaagcaaatt tcatttgaaa      300
gtgcataaat ggtcatcata agtcaaactg atcaattaga ccttcaacct aggaaacaaa      360
attttttttt tctattttaat aatacaccac actgaaatta tttgccaatg aatcccaaag      420
atttggtaca aatagtacaa ttcgtatttg ctttcctctt tcctttcttc agacaaacac      480
caaataaaat gcaggtgaaa gagatgaacc acgactagag gctgacttag aaatttatgc      540
tgactcgatc taaaaaaat tatgttggtt aatgttaatc tatctaaaat agagcatttt      600

```

```

gggaatgctt ttcaaagaag gtcaagtaac agtcatacag ctagaaaagt ccctgaaaaa 660
aagaattggt aagaagtata ataacctttt caaaacccac aatgcagctt agttttcctt 720
tatttatttg tggatcatgaa gactatcccc atttctccat aaaatcctcc ctccatactg 780
ctgcattatg gcacaaaaga ctctaagtcg caccagacag aaggaccaga gtttctgatt 840
ataacaatg atgctgggta atgtttaaat gagaacattg gatatggatg gtcag 895

```

```

<210> 74
<211> 351
<212> DNA
<213> Homo sapien

```

```

<220>
<221> misc_feature
<222> (1)...(351)
<223> n = A,T,C or G

```

```

<400> 74
tgtgcncagg ggaatgggtgg gcngtggaga ngatgacaga aaggctggaa ggaanggggg 60
tggttttgaa ggccanggcc aaggggncct caggtccgnt tctgnnaagg gacagccttg 120
aggaaggagn catggcaagc catagctagg ccaccaatca gattaagaaa nnctgagaaa 180
nctagctgac catcactggt ggtgnccagt ttcccaacac aatggaatnc caccacactg 240
gactagnnga nccactagtt ctagagcggc cgccaccgcg gtggaacccc aacttttgcc 300
cctttagnga gggttaattg cgcgcttggc ntaatcatgg tcataagctg t 351

```

```

<210> 75
<211> 251
<212> DNA
<213> Homo sapien

```

```

<400> 75
tacttgacct tctttgaaaa gcattcccaa aatgctctat tttagataga ttaacattaa 60
ccaacataat tttttttaga tcgagtcagc ataaatttct aagtcagcct ctagtctgtg 120
ttcatctctt tcacctgcat tttatttggg gtttgtctga agaaaggaaa gaggaagca 180
aatacgaatt gtactatttg taccaaactt ttgggattca ttggcaaata atttcagtgt 240
ggtgtattat t 251

```

```

<210> 76
<211> 251
<212> DNA
<213> Homo sapien

```

```

<400> 76
tatttaataa tacaccacac tgaaattatt tgccaatgaa tcccaaagat ttggtacaaa 60
tagtacaatt cgtatttgct ttctctttc ctttcttcag acaaacacca aataaaatgc 120
aggtgaaaga gatgaaccac gactagaggc tgacttagaa atttatgctg actcgatcta 180
aaaaaaatta tgttggttaa tgtaaatcta tctaaaatag agcattttgg gaatgctttt 240
caaagaaggt c 251

```

```

<210> 77
<211> 351
<212> DNA
<213> Homo sapien

```

```

<220>
<221> misc_feature

```

<222> (1) ... (351)

<223> n = A,T,C or G

<400> 77

actcaccgtg	ctgtgtgctg	tgtgcctgct	gcctggcagc	ctggccctgc	cgctgctcag	60
gaggcgggag	gcatgagtga	gctacagtgg	gaacaggctc	aggactatct	caagagannn	120
tatctctatg	actcagaaac	aaaaaatgcc	aacagttag	aagccaaact	caaggagatg	180
caaaaattct	ttggcctacc	tataactgga	atgttaaact	cccgcgcat	agaaataatg	240
cagaagccca	gatgtggagt	gccagatgtt	gcagaatact	cactatttcc	aaatagccca	300
aaatggactt	ccaaagtggg	cacctacagg	atcgatcat	atactcgaga	c	351

<210> 78

<211> 1574

<212> DNA

<213> Homo sapien

<400> 78

gccctggggg	cggaggggag	gggcccacca	cggccttatt	tccgcgagcg	ccggcactgc	60
ccgctccgag	cccggtgtctg	tcgggtgccg	agccaacttt	cctgcgtcca	tgcagccccg	120
ccggcaacgg	ctgcccgcctc	cctgggtccgg	gcccaggggc	ccgcgccccca	ccgccccgct	180
gctcgcgctg	ctgctgttgc	tcgccccggg	ggcggcgccc	gcgggggtccg	gggacccccga	240
cgaccctggg	cagcctcagg	atgctggggg	cccgcgcagg	ctcctgcagc	aggcggcgcg	300
cgcggcgctt	cactttctca	acttccgggc	cggctcgccc	agcgcgctgc	gagtgtggc	360
cgaggtgcag	gagggccgcg	cgtggattaa	tccaaaagag	ggatgtaaag	ttcacgtggt	420
cttcagcaca	gagcgctaca	acccagagtc	tttacttcag	gaaggtgagg	gacgtttggg	480
gaaatgttct	gctcgagtgt	ttttcaagaa	tcagaaaacc	agaccaacta	tcaatgtaac	540
ttgtacacgg	ctcatcgaga	aaaagaaaag	acaacaagag	gattacctgc	tttacaagca	600
aatgaagcaa	ctgaaaaacc	ccttggaat	agtcagcata	cctgataatc	atggacatat	660
tgatccctct	ctgagactca	tctgggattt	ggctttcctt	ggaagctctt	acgtgatgtg	720
ggaaatgaca	acacaggtgt	cacactacta	cttggcacag	ctcactagtg	tgaggcagtg	780
gaaaactaat	gatgatataa	ttgatattga	ttatactgtt	ctacttcatg	aattatcaac	840
acaggaaata	attccctgtc	gcattcactt	ggtctgggtac	cctggcaaac	ctcttaaagt	900
gaagtaccac	tgtcaagagc	tacagacacc	agaagaagcc	tccggaactg	aagaaggatc	960
agctgtagta	ccaacagagc	ttagtaattt	ctaaaaagaa	aaaatgatct	ttttccgact	1020
tctaaacaag	tgactatact	agcataaatc	attcttctag	taaaacagct	aaggatataga	1080
cattctaata	atttgggaaa	acctatgatt	acaagtaaaa	actcagaaat	gcaaagatgt	1140
tggttttttg	tttctcagtc	tgcttttagct	tttaactctg	gaagcgcatg	cacactgaac	1200
tctgctcagt	gctaaacagt	caccagcagg	ttcctcaggg	tttcagccct	aaaatgtaaa	1260
acctggataa	tcagtgtatg	ttgcaccaga	atcagcattt	tttttttaac	tgcaaaaaat	1320
gatggctctca	tctctgaatt	tatatctctc	attcttttga	acatactata	gctaataatat	1380
tttatgttgc	taaattgctt	ctatctagca	tgtaaataaa	agataatata	ctttcgatga	1440
aagtaaatta	taggaaaaaa	attaactgtt	ttaaaaagaa	cttgattatg	ttttatgatt	1500
tcaggcaagt	attcattttt	aacttgctac	ctacttttaa	ataaatgttt	acattttctaa	1560
aaaaaaaaaa	aaaa					1574

<210> 79

<211> 401

<212> DNA

<213> Homo sapien

<220>

<221> misc_feature

<222> (1) ... (401)

<223> n = A,T,C or G

<400> 79							
catactgtga	attgttcttg	actccttttc	ttgacattca	gttttcanaa	tttccatctt		60
tcttctggaa	ctaattgtgt	gttctcttga	ctgcctgctg	ggccagcatc	cgattgccag		120
ccagaaacgt	cacactgccc	aagatggcca	ggtacttcaa	ggtctggaac	atgttgagct		180
gagttccagt	gacatacatg	agtcccagca	tagcagcatg	tcccagggtga	aatataatcg		240
tgctaggagc	aaaagtgaag	ttggagacat	tggcaccaat	ccggatccac	tagttctaga		300
cgggccgcc	ccgcggtgga	gtctcagctt	ttgttccctt	tagtgagggt	taattgcgcg		360
cttggcqttaa	tcatggncat	agctgtttcc	tgtgtgaaat	t			401

```
<210> 80
<211> 301
<212> DNA
<213> Homo sapien
```

<400> 80						
aaaaatgaaa	catctatttt	agcagcaaga	ggctgtgagg	gatggggtag	aaaaggcatc	60
ctgagagagt	tctagaccga	cccaggtcct	gtggcacact	atacgggtca	ggaggggtgg	120
aaagacaggcc	taagctctag	gacggtgaat	ctcggggcta	tttgtggatt	tgttagaaac	180
agacattctt	ttggcctttt	cctggcactg	gtgttgcccg	caggtgggca	gaagtgagcc	240
accagtcact	gttcagtcac	tgccaccaca	gatcttcagc	agaatcttcc	ggtaatcccc	300
t						301

```
<210> 81
<211> 301
<212> DNA
<213> Homo sapien
```

```
<220>
<221> misc_feature
<222> (1)...(301)
<223> n = A,T,C or G
```

<400> 81						
tagccagggt	gctcaagcta	atttttattct	ttcccaacag	gatccatttg	gaaaatatca	60
agccttttaga	atgtggcagc	aagagaaaagc	ggactacgca	ggaacgggga	gtttgggaga	120
agctctcctg	gtgttgactt	agggatgaag	gctccaggct	gctgccagaa	atggagtcac	180
cagcagaaga	actgntttct	ctgataagga	tgtcccacca	ttttcaagct	gttcgttaaa	240
gttacacagg	tccttcttgc	agcagtaagt	accgtttagct	catttttcct	caagcggggt	300
t						301

```
<210> 82
<211> 201
<212> DNA
<213> Homo sapien
```

```
<220>
<221> misc_feature
<222> (1)...(201)
<223> n = A,T,C or G
```

<400> 82						
tcaacagaca	aaaaaagttt	attgaataca	aaactcaaag	gcatcaacag	tcttggggccc	60
aagagatcca	tggcaggaag	tcaagagttc	tgcttcaggg	tcggtctggg	cagccctgga	120
agaagtcatt	gcactatgca	gtgatgagtg	ccagggaaaac	agcatactcc	tggaaagtcc	180
acctctctqn	cactqnttca	t				201

<210> 83
<211> 251
<212> DNA
<213> Homo sapien

<220>
<221> misc_feature
<222> (1)...(251)
<223> n = A,T,C or G

<400> 83
gtaaggagca tactgtgcc atttattata gaatgcagtt aaaaaaata ttttgagggt 60
agcctctcca gtttaaaagc acttaacaag aaacacttgg acagcgatgc aatgggtctct 120
cccaaaccgg ctccctctta ccaagtaccg taaacagggt ttgagaacgt tcaatcaatt 180
tcttgatatg aacaatcaaa gcatttaatg caaacatatt tgcttctcaa anaataaaac 240
cattttccaa a 251

<210> 84
<211> 301
<212> DNA
<213> Homo sapien

<220>
<221> misc_feature
<222> (1)...(301)
<223> n = A,T,C or G

<400> 84
agttttataat gttttactat gatttagggc ttttttttca aagaacaaaa attataagca 60
taaaaactca ggtatcagaa agactcaaaa ggctgttttt cactttgttc agattttggt 120
tccaggcatt aagtgtgtca tacagttggt gccactgctg ttttccaaat gtccgatgtg 180
tgctatgact gacaactact tttctctggg tctgatcaat ttgacgtan accatttttag 240
ttcttacggc gtcnataaca aatgcttcaa catcatcagc tccaatctga agtcttgctg 300
c 301

<210> 85
<211> 201
<212> DNA
<213> Homo sapien

<400> 85
tattttgtgta tgtaacattt attgacatct acccactgca agtatagatg aataagacac 60
agtcacacca taaaggagtt tctccttaaa aggagtgaag gacattcaaa aaccaactgc 120
aataaaaaag ggtgacataa ttgctaaatg gagtggagga acagtgttca tcaattcttg 180
attgggccac aatgatatac c 201

<210> 86
<211> 301
<212> DNA
<213> Homo sapien

<220>
<221> misc_feature
<222> (1)...(301)

<223> n = A,T,C or G

<400> 86

```

tttataaaat attttattta cagtagagct ttacaaaaat agtctttaa taatacaaat    60
cccttttgca atataactta tatgactatc ttctcaaaaa cgtgacattc gattataaca    120
cataaactac atttatagtt gttaagtcac cttgtagtat aaatatgttt tcattctttt    180
tttgtaataa ggtacatacc aataacaatg aacaatggac aacaaatctt attttgntat    240
tcttccaatg taaaattcat ctctggccaa aacaaaatta accaaagaaa agtaaaacaa    300
t                                                                    301

```

<210> 87

<211> 351

<212> DNA

<213> Homo sapien

<220>

<221> misc_feature

<222> (1)...(351)

<223> n = A,T,C or G

<400> 87

```

aaaaaagatt taagatcata aataggtcat tgttgtcaca acacatttca gaatcttaaa    60
aaaacaaaca ttttggcttt ctaagaaaaa gacttttaaa aaaaatcaat tccctcatca    120
ctgaaaggac ttgtacattt ttaaacttcc agtctcctaa ggcacagtat ttaatcagaa    180
tgccaatatt accaccctgc tgtagcanga ataaagaagc aagggattaa cacttaaaaa    240
aacngccaaa ttcctgaacc aaatcattgg catthtataaa aagggataaa aaaacnggnt    300
aaggggggga gcattttaag taaagaangg ccaaggggtg tatgccngga c          351

```

<210> 88

<211> 301

<212> DNA

<213> Homo sapien

<220>

<221> misc_feature

<222> (1)...(301)

<223> n = A,T,C or G

<400> 88

```

gttttaggtc ttaccaatt tgattggttt atcaacaggg catgaggttt aaatatatct    60
ttgaggaaag gtaaagtcaa atttgacttc ataggtcatc ggcgtcctca ctctgtgca    120
ttttctggtg gaagcacaca gttaattaac tcaagtgtgg cgntagegat gctttttcat    180
ggngtcattt atccacttgg tgaacttgca cacttgaatg naaactcctg ggtcattggg    240
ntggccgcaa gggaaagggtc cccaagacac caaaccttgc agggtagctn tgcacaccaa    300
c                                                                    301

```

<210> 89

<211> 591

<212> DNA

<213> Homo sapien

<400> 89

```

tttttttttt tttttttatt aatcaaataa ttcaaaacaa ccatcattct gtcaatgccc    60
aagaccccag ctggtcctct cccacatgt cacactctcc tcagcctctc ccccaaccct    120
gctctccctc ctcccctgcc ctagcccagg gacagagtct agggaggagc tggggcagag    180

```

ctggaggcag	gaagagagca	ctggacagac	agctatgggt	tggattgggg	aagagattag	240
gaagtaggtt	cttaagacc	cttttttagt	accagatata	cagccatatt	cccagctcca	300
ttattcaaat	catttcccat	agcccagctc	ctctctgttc	tccccctact	accaattctt	360
tggctcttac	acaattttta	tccctcaa	attcatccct	ggcccaacca	gtcccctgag	420
cctccctctg	gtggagactc	ctccacccat	gagctcccca	gagcatccaa	gacagagtgc	480
acagagacct	ggggaaggaa	gctgaacttt	gcagagatgt	ggacaggtgc	aggctagggt	540
acagggtggt	ggtagaggag	acaagtttta	tttccaggcc	cacagtctct	c	591

<210> 90

<211> 1978

<212> DNA

<213> Homo sapien

<400> 90

tttttttttt	ttttttatca	aatgaatact	ttattagaga	cataacacgt	ataaaataaa	60
tttcttttca	tcattggagt	accagatttt	aaaaccaacc	aacactttct	cattttttaca	120
gctaagacat	gttaaatctt	taaatgccat	aattttttgt	caactgcttt	gtcattcaac	180
tcacaagtct	agaatgtgat	taagctacaa	atctaagtat	tcacagatgt	gtcttaggct	240
tggtttgtaa	caatctagaa	gcaatctgtt	tacaaaagt	ccaccaaaagc	attttaaaga	300
aaccaattta	atgccaccaa	acataagcct	gctataacct	ggaaacaaaa	aatctcacac	360
ctaaattcta	gcagagtaaa	cgattccaac	tagaatgtac	tgtatatcca	tatggcacat	420
ttatgacttt	gtaatatgta	attcataata	cagggtttagg	tgtgtggtat	ggagctagga	480
aaaccaaagt	agtaggatat	tatagaaaag	atctgatgtt	aagtataaag	tcatatgcct	540
gatttctctca	aaccttttgt	ttttctctcat	gtcttctgtc	tttatatttt	tatcacaac	600
caagatctaa	cagggttctt	tctagaggat	tattagataa	gtaacacttg	atcattaagc	660
acggatcatg	ccactcattc	atggttggtc	tatgttccat	gaactcta	agcccaactt	720
atacatggca	ctccaagggg	atgcttcagc	cagaaagtaa	agggctgaaa	aagtagaaca	780
atacaaaagc	cctcgtgtgg	tgggaactgt	ggcctcactc	ttacttgtec	ttccattcaa	840
aacagtttgg	cacctttcca	tgacgaggat	ctctacaggt	aggttaaaat	acttttctgt	900
gctattcagc	cagaaatagt	ttttgtgctg	gatatgattt	taaaacagat	tttgtctgtc	960
accagtgcga	aaacattaca	gatgtctggg	ctaatacaaa	aacacataag	aatctacaac	1020
tttatattta	atactctatt	caaatttaac	tcaaagta	gcaaaataat	tagaagtaaa	1080
aacttaattc	ttctgagagc	tctatttgga	aaagcttcac	atatccacac	acaaatatgg	1140
gtatattcat	gcacagggca	aacaactgta	ttctgaagca	taaataaact	caaagtaaga	1200
catcagtagc	tagataccag	ttccagtatt	ggttaatggg	ctctggggat	cccattttaa	1260
gcactctcag	atgaggatct	tgctcagttg	ttagactatc	attagtttga	ttaagcaact	1320
gaagtttact	tcataaatta	ctttttccta	tatccaggac	tctgcctgag	aaattttata	1380
cattcctcca	aaggtaagta	ttctccaaag	gtaagtattt	gactattaac	acaaaggcaa	1440
tgtgattatt	gcataatgac	actaaatat	atgtggcttt	tctgttaggt	ttataagttt	1500
tcaatgatca	gttcaagaaa	atgcagatca	tatataacta	aggttttaca	ccagtggttg	1560
acaaactatg	gcccacaggc	taaaccagc	ctccccctgt	ttttataaat	aagttttatt	1620
agacataacc	acactcattc	atttctgtat	tgtgtatagc	tgctttcacg	ctatactagc	1680
agaactgaat	agttgtgaca	gagactgtat	ggaccgtgaa	gcataaatat	ttaccatctg	1740
gcccattcta	aaaaaagtgt	gccaatccct	ggtttacact	aaaatataga	gtttagtggg	1800
aagcctattt	gaaatgtgtt	tttttttagg	gctgtaatta	ccaattaaaa	ttaaggttca	1860
ggtgactcag	caaccaaaca	aaagggatac	taatttttta	tgaacaatat	atttgtattt	1920
tatggacata	aaaggaaact	ttcagaaaga	aaaggaggaa	aataaagggg	gaaaggga	1978

<210> 91

<211> 895

<212> DNA

<213> Homo sapien

<400> 91

tttttttttt	ttttttcttg	tttaaaaaaa	ttgttttcat	tttaatgata	tgagttagta	60
------------	------------	------------	------------	------------	------------	----

acaaacaaat	gtacaaaatt	gtctttcaca	tttccatata	ttgtgttatg	gaccaaata	120
aaacgctgga	ctacaaatgc	agggtttctt	atatccttaa	cttcaattat	tgctacttat	180
aaataaaagg	gatttgctaa	cacatgcatt	tgtgaacaca	gatgccaaaa	attatacatg	240
taagttaatg	cacaaccaag	agtatacact	gttcatttgt	gcagttatgc	gtcaaagcg	300
actgacacag	aagcagttat	cctgggatat	ttcactctat	atgaaaagca	tcttggagaa	360
atagattgaa	atacagttta	aaacaaaaat	tgtattctac	aaatacaata	aaatttgcaa	420
cttgacatc	tgaagcaaca	tttgagaaag	ctgcttcaat	aacctgctg	ttatatgggt	480
tttataggta	tatctccaaa	gtcatgggtt	gggatatagc	tgctttaaag	aaaataaata	540
tgtatattaa	aaggaaaatc	acactttaaa	aatgtgagga	aagctttgaa	aacagtctta	600
atgcatgagt	ccatctacat	attttcaagt	tttggaacaa	gaaagaagtt	tagaattttc	660
aaagtaatct	gaaaactttc	taagccattt	taaaataaga	tttttttccc	catctttcca	720
atgtttccta	tttgatagtg	taatacagaa	atgggcagtt	tctagtgtca	acttaactgt	780
gctaattcat	aagtcattat	acatttatga	cttaagagtt	caaataagtg	gaaattgggt	840
tataatgaaa	atgacaaggg	ggcccttca	gcagccactc	atctgaacta	gtaat	895

<210> 92

<211> 1692

<212> DNA

<213> Homo sapien

<400> 92

tttttttttt	tttttaactt	ttagcagtg	ttatttttgt	taaaagaaac	caattgaatt	60
gaaggccaag	acaccttctg	attgcacaga	ttaaacaaga	aagtattact	tatttcaact	120
ttacaaagca	tcttattgat	ttaaaaagat	ccatactatt	gataaagttc	accatgaaca	180
tatatgtaat	aaggagacta	aaatattcat	tttaccatct	tacaacatgt	atttcatatt	240
tctaatacaac	cacaaatcat	ataggaaaat	atttaggtcc	atgaaaaagt	ttcaaaacat	300
taaaaaatta	aagttttgaa	acaaatcaca	tgtgaaagct	cattaaataa	taacattgac	360
aaataaatag	ttaatcagct	ttacttatta	gctgctgcca	tgcatctctg	gcattccatt	420
ccaagcgagg	gtcagcatgc	agggtataat	ttcatactat	gcgaccgtaa	agagctacag	480
ggcttatttt	tgaagtgaag	tgtcacaggg	tctttcattc	tctttcaaaag	gaagatcact	540
catggctgct	aaactgttcc	catgaagagt	acaaaaaaag	cacctttctg	aaatgttact	600
gtgaagattc	atgacaacat	atttttttta	acctgttttg	aaggagtttt	gtttaggaga	660
ggggatgggc	cagtagatgg	agggtatctg	agaagccctt	ttctgtttta	aaatataatg	720
attcactgat	gtttatagta	tcaacagctc	tttaagaaca	atgaggaatt	aaaactacag	780
gatacgtgga	atttaaatgc	aaattgcatt	catggatata	cctacatctt	gaaaaacttg	840
aaaaggaaaa	actattccca	aagaagggtcc	tgatacttaa	gacagcttgc	tgggtttgat	900
caaagcagaa	agcatatact	ttcaagttag	aaaacagcag	tggcaggctt	gagtcctcca	960
agcaatcaaa	tctgtaaagc	agatgggttac	tagtaagtct	agttatggga	gtctgagttc	1020
taactcatgc	tgtgcttgct	ggatttgctg	gctcttttcc	gctctctgtg	atgctggact	1080
ggcttgctcag	gtgacatgct	ctcaaagtgt	tgactggact	cgttggtgctg	ccgggtgtac	1140
ctcttgcact	tgaggcagct	gactactgtg	attttgtagg	tgcgtgtgct	gccatcttgg	1200
cactgcagct	ggattctctg	ggtaggggtt	ttgtcattga	cacaccgcca	ctcctgggag	1260
ctcctcctgc	tccagtactt	tgttccatag	cctcctccaa	tccagttagg	gagcactggc	1320
aggggcaagc	actcgccagc	acacaccagc	tccttcagag	ggctgatgct	ggtgcactgg	1380
ccatcagaga	tgtatttggt	ggaacgcagt	tcccggcaac	ccacttgaac	ccgagtgttc	1440
cgatccagtc	cagtgttact	gaaatgcctg	cctccatttc	tggcttgatt	caacgtgctg	1500
ttgctgctgg	ggtgtgctgg	aacaggttta	accacatgtg	aataaaggat	ttctgtggca	1560
tcatttttaa	aagccaaaca	gcttttcatt	aggatgcatg	caaggggaag	gagatagaaa	1620
tgaatggcag	gaggaagcat	ggtgagtaga	ggatttgctt	gactgaagag	ctgggttaatt	1680
cttttgcttc	tg					1692

<210> 93

<211> 251

<212> DNA

<213> Homo sapien

<400> 93

```

ccccacctac ccaaattatta gacaccaaca cagaaaagct agcaatggat tcccttctac      60
tttgtaaata aaataagtta aatatttaaa tgcctgtgtc tctgtgatgg caacagaagg      120
accaaagggc cacatcctga taaaaggtaa gaggggggtg gatcagcaaa aagacagtgc      180
tgtgggctga ggggacctgg ttcttgtgtg ttgccccctca agactcttcc cctacaaata      240
actttcatat g                                     251

```

<210> 94

<211> 735

<212> DNA

<213> Homo sapien

<400> 94

```

tttttttttt tttttccact tctcagttta tttctgggac taaatttggg tcagagctgc      60
agagaaggga tgggccctga gcttgaggat gaaagtgtcc cagggagatt gagacgcaac      120
ccccgccctg gacagttttg gaaattgttc ccagggttca actagagaga cacggtcagc      180
ccaatgtggg ggaagcagac cctgagtcca ggagacatgg ggtcaggggc tggagagatg      240
aacatttcca acatctctgg gaaggaaatga gggctctgaaa ggagtgtcag ggctgtccct      300
gcagcaggtg gggatgccgg tgtgctgagt cctgggatga ctcaggagtt ggcctggatg      360
gtttcctgga tccacttggg gaacttgtag aggttcgtgt agacacccgg tctgttgggc      420
cgggcacaag ggtaatctcc ccaggacacg agtccctgca gggagccatt gcagaccaca      480
ggccccccag aatcacctcg gcaggagtct ctacctgctt tgtcaccggc gcagaacatg      540
gtgtcatcta tctgtctcgg gtaagcatcc tcgcaccttt tctgacttag cacgctgata      600
ttcaagcaact ggaggacctt agggaaagtgc acttgggggc tcttgggtgt cccccagcca      660
gacaccaagc actttgtccc agcagaggga caatgagagg agacgttgat gggctctgaca      720
tcttttagtg gacga                                     735

```

<210> 95

<211> 578

<212> DNA

<213> Homo sapien

<400> 95

```

cttgccctct cttaggcttt gaagcatttt tgtctgtgct ccctgatctt caggtcacca      60
ccatgaagtt cttagcagtc ctggtactct tgggagtttc catctttctg gtctctgccc      120
agaatccgac aacagctgct ccagctgaca cgtatccagc tactggctct gctgatgatg      180
aagccctga tgctgaaacc actgctgctg caaccactgc gaccactgct gctcctacca      240
ctgcaaccac cgctgcttct accactgctc gtaaagacat tccagtttta cccaaatggg      300
ttggggatct cccgaatggg agagtgtgtc cctgagatgg aatcagcttg agtcttctgc      360
aattgggtcac aactattcat gcttctgtgt atttcatcca actacttacc ttgcctacga      420
tatccccctt atctctaata agtttatttt ctttcaaata aaaaataact atgagcaaca      480
aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa      540
aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa                                     578

```

<210> 96

<211> 594

<212> DNA

<213> Homo sapien

<400> 96

```

atggcaaaga atggacttgt aatttgcata ctggtgatca ccttactcct ggaccagacc      60
accagccaca catccagatt aaaagccagg aagcacagca aacgtcgagt gagagacaag      120
gatggagatc tgaagactca aattgaaaag ctctggacag aagtcaatgc cttgaaggaa      180
attcaagccc tgcagacagt ctgtctccga ggcactaaag ttcacaagaa atgctacctt      240

```

gcttcagaag	gtttgaagca	tttccatgag	gccaatgaag	actgcatttc	caaaggagga	300
atcctgggta	tccccaggaa	ctccgacgaa	atcaacgccc	tccaagacta	tggtaaaagg	360
agcctgccag	gtgtcaatga	cttttggtcg	ggcatcaatg	acatggtcac	ggaaggcaag	420
tttgttgacg	tcaacggaat	cgctatctcc	ttcctcaact	gggaccgtgc	acagcctaac	480
gggtggcaagc	gagaaaactg	tgtcctgttc	tcccaatcag	ctcagggcaa	gtggagtgat	540
gaggcctgtc	gcagcagcaa	gagatacata	tgcgagttca	ccatccctca	atag	594

<210> 97

<211> 3101

<212> DNA

<213> Homo sapien

<400> 97

tgttggggcc	tcagcctccc	aagtagctgg	gactacaggt	gcctgccacc	acgccagct	60
aatTTTTTgt	atatTTTTta	gtagagacgg	ggtttcaccg	tggctcfaat	ctcctgacct	120
cgtgatctgc	cagccttggc	ctcccaaagt	gtattctctt	tttattatta	ttattatttt	180
tgagatggag	tctgtctctg	tcgcccaggc	tggagtgcag	tgggtcgatc	tctgtcact	240
gcaagctccg	cctcctgggt	tcatgccatt	ctcctgcctc	agcctcccga	gtagctggga	300
ctacaggccc	ctgccaccac	acccggctaa	ttttttgtat	tttttagtaga	gacaggggtt	360
caccatgtta	gccagggtgg	tctctatctt	ctgacctcgt	gatccgcctg	cctcagttct	420
tcaaagtgtc	gggattacag	gcgtgagcca	ccgcgaccag	ccaactattg	ctgtttattt	480
ttaaatatat	tttaaagaaa	caattagatt	tgttttcttt	ctcattcttt	tacttctact	540
cttcatgtat	gtataattat	atttgtgttt	tctattacct	tttctccttt	tactgtattg	600
gactataata	attgtgtctc	ctaatttctg	ttcactaata	ttatcagctt	agataaatac	660
ttaatTTTTa	acttatatat	tgagtattaa	attgatcagt	tttatttgta	attatctatc	720
ttccgcttgg	ctgaatataa	cttcttaagc	ttataacttc	ttgttctttc	catgttattt	780
ttttcttttt	tttaattgtat	tgaatttctt	ctgacactca	ttctagtaac	ttttttctcg	840
gtgtgcaacg	taagttataa	tttgtttctc	agatttgaga	tctgccataa	gtttgaggct	900
ttatTTTTtt	tttttatttg	ctttatggca	agtcggacaa	cctgcatgga	tttggcatca	960
atgtagtcac	ccatatctaa	gagcagcact	tgcttcttag	catgatgagt	tgtttctgga	1020
ttgtttcttt	attttactta	tattcctggg	agattcttat	attttccctt	caactctatt	1080
cagcatttta	ggaattctta	ggactttctg	agaatttttag	ctttctgtat	taaatgtttt	1140
taatgagtat	tgcattttct	caaaaagcac	aaatatcaat	agtgtacaca	tgaggaaaac	1200
tatatatata	ttctgttgca	gatgacagca	tctcataaca	aaatcctagt	tacttcattt	1260
aaaagacagc	tctcctccaa	tatactatga	ggtaacaaaa	atttgtagtg	tgtaattttt	1320
ttaatattag	aaaactcatc	ttacattgtg	cacaaatttc	tgaagtata	atacttcact	1380
gtttttctat	agaagtaact	taatatggc	aaaattactt	atttgaattt	aggttttggc	1440
tttcatcata	tacttctctc	ttaacatttc	cctcaatcca	taaatgcaat	ctcagtttga	1500
atcttccatt	taaccagaa	gttaattttt	aaaaccttaa	taaaatttga	atgtagctag	1560
atattatttg	ttggttacat	attagtcfaat	aatttatatt	acttacaatg	atcagaaaaat	1620
atgatctgaa	tttctgtctg	cataaattca	ataacgtatt	ttaggcctaa	acctttccat	1680
ttcaaactct	tgggtctggg	aattgaaaat	aatcattatc	ttttgttttc	tggccaaaaa	1740
tgtcgcccat	ttatttctat	ccctaattag	tcaaactttc	taataaatgt	atttaacgtt	1800
aatgatgttt	atttgcttgt	tgtatactaa	aaccattagt	ttctataatt	taaatgtcac	1860
ctaatatgag	tgaaaatgtg	tcagaggctg	gggaagaatg	tggatggaga	aagggaaggt	1920
gttgatcaaa	aagtacccaa	gtttcagtta	cacaggaggc	atgagattga	tctagtgcac	1980
aaaatgatga	gtataataaa	taataatgca	ctgtatattt	tgaatttgct	aaaagtagat	2040
ttaaaattga	tttacataat	attttacata	tttataaagc	acatgcaata	tgttgtttaca	2100
tgtatagaat	gtgcaacgat	caagtcaggg	tatctgtggg	atccaccact	ttgagcattt	2160
atcgattcta	tatgtcagga	acatttcaag	ttatctgttc	tagcaaggaa	atataaaata	2220
cattatagtt	aactatggcc	tatctacagt	gcaactaaac	actagatttt	attcctttcc	2280
aactgtgggt	ttgtattcat	ttaccacctt	cttttcattc	cctttctcac	ccacacactg	2340
tgccgggccc	caggcatata	ctattctact	gtctgtctct	gtaaggatta	tcatttttagc	2400
ttccacatat	gagagaatgc	atgcaaagtt	tttctttcca	tgtctggctt	atttcactta	2460
acaaaatgac	ctccgcttcc	atccatgtta	tttatattac	ccaatagtgt	tcataaatat	2520

```

atatacacac atatatacca cattgcattt gtccaattat tcattgacgg aaactgggta 2580
atgttatatc gttgctattg tgaatagtgc tgcaataaac acgcaagtgg ggatataatt 2640
tgaagagttt ttttgttgat gttccataca aattttaaga ttgttttgtc tatgtttgtg 2700
aaaatggcgt tagtattttc atagagattg cattgaatct gtagattgct ttgggtaagt 2760
atggttattt tgatgggtatt aattttttca ttccatgaag atgagatgct ttccatttg 2820
tttgtgtcct ctacattttc tttcatcaaa gttttgttgt atttttgaag tagatgtatt 2880
tcaccttata gatcaagtgt attccctaaa tattttattt ttgtagctat tgtagatgaa 2940
attgccttct cgattttctt ttcacttaat tcattattag tgtatggaaa tgttatggat 3000
ttttatttgt tgggttttta tcaaaaactg tattaactt agagttttt gtggagttt 3060
taagtttttc tagatataag atcatgacat ctacaaaaa a 3101

```

<210> 98
 <211> 90
 <212> PRT
 <213> Homo sapien

<400> 98

```

Met Lys Phe Leu Ala Val Leu Val Leu Leu Gly Val Ser Ile Phe Leu
 1          5          10          15
Val Ser Ala Gln Asn Pro Thr Thr Ala Pro Ala Asp Thr Tyr Pro
          20          25          30
Ala Thr Gly Pro Ala Asp Asp Glu Ala Pro Asp Ala Glu Thr Thr Ala
          35          40          45
Ala Ala Thr Thr Ala Thr Thr Ala Ala Pro Thr Thr Ala Thr Thr Ala
          50          55          60
Ala Ser Thr Thr Ala Arg Lys Asp Ile Pro Val Leu Pro Lys Trp Val
65          70          75          80
Gly Asp Leu Pro Asn Gly Arg Val Cys Pro
          85          90

```

<210> 99
 <211> 197
 <212> PRT
 <213> Homo sapien

<400> 99

```

Met Ala Lys Asn Gly Leu Val Ile Cys Ile Leu Val Ile Thr Leu Leu
 1          5          10          15
Leu Asp Gln Thr Thr Ser His Thr Ser Arg Leu Lys Ala Arg Lys His
          20          25          30
Ser Lys Arg Arg Val Arg Asp Lys Asp Gly Asp Leu Lys Thr Gln Ile
          35          40          45
Glu Lys Leu Trp Thr Glu Val Asn Ala Leu Lys Glu Ile Gln Ala Leu
          50          55          60
Gln Thr Val Cys Leu Arg Gly Thr Lys Val His Lys Lys Cys Tyr Leu
65          70          75          80
Ala Ser Glu Gly Leu Lys His Phe His Glu Ala Asn Glu Asp Cys Ile
          85          90          95
Ser Lys Gly Gly Ile Leu Val Ile Pro Arg Asn Ser Asp Glu Ile Asn
          100          105          110
Ala Leu Gln Asp Tyr Gly Lys Arg Ser Leu Pro Gly Val Asn Asp Phe
          115          120          125
Trp Leu Gly Ile Asn Asp Met Val Thr Glu Gly Lys Phe Val Asp Val
          130          135          140
Asn Gly Ile Ala Ile Ser Phe Leu Asn Trp Asp Arg Ala Gln Pro Asn

```

145 150 155 160
 Gly Gly Lys Arg Glu Asn Cys Val Leu Phe Ser Gln Ser Ala Gln Gly
 165 170 175
 Lys Trp Ser Asp Glu Ala Cys Arg Ser Ser Lys Arg Tyr Ile Cys Glu
 180 185 190
 Phe Thr Ile Pro Gln
 195

<210> 100
 <211> 3410
 <212> DNA
 <213> Homo sapien

<400> 100
 gggaaccagc ctgcacgcgc tggctccggg tgacagccgc gcgcctcggc caggatctga 60
 gtgatgagac gtgtcccccac tgaggtgccc cacagcagca ggtgttgagc atgggctgag 120
 aagctggacc ggcaccaaag ggctggcaga aatgggcgcc tggctgattc ctaggcagtt 180
 ggccgagca aggaggagag gccgcagctt ctggagcaga gccgagacga agcagttctg 240
 gagtgcctga acggccccct gagccctacc cgcctggccc actatggtcc agaggctgtg 300
 ggtgagccgc ctgtgcgggc accggaagc ccagctcttg ctggtcaacc tgctaaccctt 360
 tggcctggag gtgtgttttg ccgcaggcat cacctatgtg ccgcctctgc tgcctggaagt 420
 gggggtagag gagaagttca tgaccatggt gctgggcatt ggtccagtgc tgggcctggt 480
 ctgtgtcccg ctctagggt cagccagtga ccactggcgt ggacgctatg gccgccgccg 540
 gcccttcacg tgggcaactgt ccttgggcat cctgctgagc ctctttctca tcccaagggc 600
 cggctggcta gcagggtgc tgtgcccga tcccaggccc ctggagctgg cactgctcat 660
 cctgggcgtg gggctgctgg acttctgtgg ccagggtgtg ttcactccac tggaggccct 720
 gctctctgac ctcttcgggg acccggacca ctgtcgccag gcctactctg tctatgcctt 780
 catgatcagt cttgggggct gcctgggcta cctcctgcct gccattgact gggacaccag 840
 tgccctggcc ccctacctgg gcaccagga ggagtgcctc tttggcctgc tcacctcat 900
 ctctctcacc tgcgtagcag ccacactgct ggtggctgag gaggcagcgc tgggccccac 960
 cgagccagca gaagggtgt cggccccctc ctgtcgccc cactgctgtc catgccgggc 1020
 ccgcttggtt ttcgggaacc tgggcgccct gcttccccgg ctgcaccagc tgtgctgccg 1080
 catgccccgc accctgcgcc ggctcttcgt ggtgagctg tgcagctgga tggcactcat 1140
 gaccttcacg ctgttttaca cggatttcgt gggcgagggg ctgtaccagg gcgtgcccag 1200
 agctgagccg ggcaccgagg cccggagaca ctatgatgaa ggcgttcgga tgggcagcct 1260
 ggggctgttc ctgcagtgcg ccactctcct ggtcttctct ctggctcatg accggctggt 1320
 gcagcgattc ggcactcgag cagtctattt ggccagtgtg gcagctttcc ctgtggctgc 1380
 cggtgccaca tgcctgtccc acagtgtggc cgtggtgaca gcttcagccg ccctcaccgg 1440
 gtacaccttc tcagccctgc agatcctgcc ctacacactg gcctccctct accaccggga 1500
 gaagcaggtg ttctgcccc aataccgagg ggacactgga ggtgctagca gtgaggacag 1560
 cctgatgacc agcttccctgc caggccctaa gcctggagct ccctcccta atggacacgt 1620
 tgggtgctgga ggcagtggcc tgcctccacc tccaccgcg ctctgcgggg cctctgcctg 1680
 tgatgtctcc gtacgtgtgg tgggtgggtga gcccaccgag gccagggtgg ttccggggccg 1740
 gggcatctgc ctggacctgc ccactctgga tagtgcttc ctgctgtccc aggtggcccc 1800
 atccctgttt atgggtcca ttgtccagct cagccagtct gtcactgcct atatggtgtc 1860
 tgccgcaggc ctgggtctgg tcgccattta ctttgttaca caggtagtat ttgacaagag 1920
 cgacttggcc aaatactcag cgtagaaaac ttccagcaca ttgggggtgga gggcctgcct 1980
 cactgggtcc cagctcccc ctctgttag ccccatgggg ctgccgggct ggccgccagt 2040
 ttctgttgct gccaaagtaa tgtggctctc tgcctgccacc ctgtgctgct gaggtgcgta 2100
 gctgcacagc tgggggctgg ggcgtccctc tcctctctcc ccagtctcta gggtgcctg 2160
 actggaggcc ttccaagggg gtttcagtct ggacttatac agggaggcca gaagggtcc 2220
 atgcaactga atgcggggac tctgcagggt gattaccag gctcagggtt aacagctagc 2280
 ctctagtgtg agacacacct agagaagggt ttttgggagc tgaataaact cagtcacctg 2340
 gtttcccatc tctaagcccc ttaacctgca gcttcgttta atgtagctct tgcattgggag 2400
 tttctaggat gaaacactcc tccatgggat ttgaacatat gacttatttg taggggaaga 2460

```

gtcctgaggg gcaacacaca agaaccaggt cccctcagcc cacagcactg tctttttgct 2520
gatccacccc cctcttacct tttatcagga tgtggcctgt tggccttct gttgccatca 2580
cagagacaca ggcatttaaa tatttaactt atttatTTaa caaagtagaa gggaaatccat 2640
tgctagcttt tctgtgttgg tgtctaatat ttgggtaggg tgggggatcc ccaacaatca 2700
ggccccctga gatagctggg cattgggctg atcattgcca gaatcttctt ctcctggggg 2760
ctggccccc aaatgccta acccaggacc ttggaaattc tactcatccc aaatgataat 2820
tccaaatgct gttacccaag gttaggggtg tgaaggaagg tagagggtag ggcttcagggt 2880
ctcaacggct tccctaacca cccctcttct cttggccag cctggttccc cccacttcca 2940
ctccccctta ctctctctag gactgggctg atgaaggcac tgcccaaaat ttccccctacc 3000
cccaactttc cctaccccc aactttcccc accagctcca caaccctgtt tggagctact 3060
gcaggaccag aagcacaag tgcggtttcc caagcctttg tccatctcag ccccagagt 3120
atatctgtgc ttgggggaatc tcacacagaa actcaggagc accccctgcc tgagctaagg 3180
gaggtcttat ctctcagggg ggggtttaagt gccgtttgca ataatgtcgt cttattttatt 3240
tagcgggggtg aatattttat actgtaagtg agcaatcaga gtataatgtt tatggtgaca 3300
aaattaaagg ctttcttata tgtttaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 3360
aaaaaaaaara aaaaaaaaaa aaaaaaaaaa aaaaaataaa aaaaaaaaaa 3410

```

<210> 101

<211> 553

<212> PRT

<213> Homo sapien

<400> 101

```

Met Val Gln Arg Leu Trp Val Ser Arg Leu Leu Arg His Arg Lys Ala
1          5          10          15
Gln Leu Leu Leu Val Asn Leu Leu Thr Phe Gly Leu Glu Val Cys Leu
20          25          30
Ala Ala Gly Ile Thr Tyr Val Pro Leu Leu Leu Glu Val Gly Val
35          40          45
Glu Glu Lys Phe Met Thr Met Val Leu Gly Ile Gly Pro Val Leu Gly
50          55          60
Leu Val Cys Val Pro Leu Leu Gly Ser Ala Ser Asp His Trp Arg Gly
65          70          75          80
Arg Tyr Gly Arg Arg Arg Pro Phe Ile Trp Ala Leu Ser Leu Gly Ile
85          90          95
Leu Leu Ser Leu Phe Leu Ile Pro Arg Ala Gly Trp Leu Ala Gly Leu
100         105         110
Leu Cys Pro Asp Pro Arg Pro Leu Glu Leu Ala Leu Leu Ile Leu Gly
115         120         125
Val Gly Leu Leu Asp Phe Cys Gly Gln Val Cys Phe Thr Pro Leu Glu
130         135         140
Ala Leu Leu Ser Asp Leu Phe Arg Asp Pro Asp His Cys Arg Gln Ala
145         150         155         160
Tyr Ser Val Tyr Ala Phe Met Ile Ser Leu Gly Gly Cys Leu Gly Tyr
165         170         175
Leu Leu Pro Ala Ile Asp Trp Asp Thr Ser Ala Leu Ala Pro Tyr Leu
180         185         190
Gly Thr Gln Glu Glu Cys Leu Phe Gly Leu Leu Thr Leu Ile Phe Leu
195         200         205
Thr Cys Val Ala Ala Thr Leu Val Ala Glu Glu Ala Ala Leu Gly
210         215         220
Pro Thr Glu Pro Ala Glu Gly Leu Ser Ala Pro Ser Leu Ser Pro His
225         230         235         240
Cys Cys Pro Cys Arg Ala Arg Leu Ala Phe Arg Asn Leu Gly Ala Leu
245         250         255

```


Leu Pro Arg Leu His Gln Leu Cys Cys Arg Met Pro Arg Thr Leu Arg
 260 265 270
 Arg Leu Phe Val Ala Glu Leu Cys Ser Trp Met Ala Leu Met Thr Phe
 275 280 285
 Thr Leu Phe Tyr Thr Asp Phe Val Gly Glu Gly Leu Tyr Gln Gly Val
 290 295 300
 Pro Arg Ala Glu Pro Gly Thr Glu Ala Arg Arg His Tyr Asp Glu Gly
 305 310 315 320
 Val Arg Met Gly Ser Leu Gly Leu Phe Leu Gln Cys Ala Ile Ser Leu
 325 330 335
 Val Phe Ser Leu Val Met Asp Arg Leu Val Gln Arg Phe Gly Thr Arg
 340 345 350
 Ala Val Tyr Leu Ala Ser Val Ala Ala Phe Pro Val Ala Ala Gly Ala
 355 360 365
 Thr Cys Leu Ser His Ser Val Ala Val Val Thr Ala Ser Ala Ala Leu
 370 375 380
 Thr Gly Phe Thr Phe Ser Ala Leu Gln Ile Leu Pro Tyr Thr Leu Ala
 385 390 395 400
 Ser Leu Tyr His Arg Glu Lys Gln Val Phe Leu Pro Lys Tyr Arg Gly
 405 410 415
 Asp Thr Gly Gly Ala Ser Ser Glu Asp Ser Leu Met Thr Ser Phe Leu
 420 425 430
 Pro Gly Pro Lys Pro Gly Ala Pro Phe Pro Asn Gly His Val Gly Ala
 435 440 445
 Gly Gly Ser Gly Leu Leu Pro Pro Pro Pro Ala Leu Cys Gly Ala Ser
 450 455 460
 Ala Cys Asp Val Ser Val Arg Val Val Val Gly Glu Pro Thr Glu Ala
 465 470 475 480
 Arg Val Val Pro Gly Arg Gly Ile Cys Leu Asp Leu Ala Ile Leu Asp
 485 490 495
 Ser Ala Phe Leu Leu Ser Gln Val Ala Pro Ser Leu Phe Met Gly Ser
 500 505 510
 Ile Val Gln Leu Ser Gln Ser Val Thr Ala Tyr Met Val Ser Ala Ala
 515 520 525
 Gly Leu Gly Leu Val Ala Ile Tyr Phe Ala Thr Gln Val Val Phe Asp
 530 535 540
 Lys Ser Asp Leu Ala Lys Tyr Ser Ala
 545 550

<210> 102

<211> 940

<212> DNA

<213> Human

<400> 102

tttactgctt	ggcaaagtac	cctgagcatc	agcagagatg	ccgagatgaa	atcaggggaac	60
tcctagggga	tgggtcttct	attacctggg	aacacctgag	ccagatgcct	tacaccacga	120
tgtgcatcaa	ggaatgcctc	cgctctacg	caccggtagt	aaacatatcc	cggttactcg	180
acaaacccat	cacctttcca	gatggacgct	ccttacctgc	aggaataact	gtgtttatca	240
atatttgggc	tcttcaccac	aacccctatt	tctgggaaga	ccctcagggtc	tttaacccct	300
tgagattctc	cagggaaaat	tctgaaaaaa	tacatcccta	tgccctcata	ccattctcag	360
ctggattaag	gaactgcatt	gggcagcatt	ttgccataat	tgagtgtaaa	gtggcagtgg	420
cattaactct	gtccgccttc	aagctggctc	cagaccactc	aaggcctccc	cagcctgttc	480
gtcaagtgtg	cctcaagtcc	aagaatggaa	tccatgtgtt	tgcaaaaaaa	gtttgctaata	540
tttaagtcct	ttcgtataag	aattaatgag	acaattttcc	taccaaagga	agaacaaaag	600

gataaatata	atacaaaata	tatgtatatg	gttggttgac	aaattatata	acttaggata	660
cttctgactg	gttttgacat	ccattaacag	taattttaat	ttctttgctg	tatctgggtga	720
aaccacacaa	aacmctgaa	aaaactcaag	ctgacttcca	ctgcgaagg	aaattattgg	780
tttggtgaac	tagtggtaga	gtggccttca	agcatagttt	gatcaaaact	ccactcagta	840
tctgcattac	ttttatyt	gcaaatact	gcatgatagc	tttattytca	gttatctttc	900
cccataataa	aaaatatctg	ccaaaaaaaa	aaaaaaaaaa			940

<210> 103

<211> 529

<212> DNA

<213> Human

<400> 103

tttttttttt	tttttactga	tagatggaat	ttattaagct	tttcacatgt	gatagcacat	60
agttttaatt	gcatccaaag	tactaacaaa	aactctagca	atcaaraatg	gcagcatgtt	120
attttataac	aatcaacacc	tgtggccttt	aaaatttgg	tttcataara	taatttatac	180
tgaagtaaat	ctagccatgc	ttttaaaaaa	tgcttttaggt	cactccaagc	ttggcagtta	240
acatttgcca	taaacaataa	taaaacaatc	acaatttaat	aaataacaaa	tacaacattg	300
taggccataa	tcatatacag	tataaggaaa	aggkggtagt	gttgagtaag	cagttattag	360
aatagaatac	cttgccctct	atgcaaatat	gtctaracac	tttgattcac	tcagccctga	420
cattcagttt	tcaaagtagg	agacagggtc	tacagtatca	ttttacagtt	tccaacacat	480
tgaaaacaag	tagaaaatga	tgagttgatt	tttattaatg	cattacatc		529

<210> 104

<211> 469

<212> DNA

<213> Human

<400> 104

cccaacacaa	tggataaaaa	cacttatagt	aaatggggac	attcactata	atgatctaag	60
aagctacaga	ttgtcatagt	tgttttcctg	ctttacaaaa	ttgtccaga	tctggaatgc	120
cagtttgacc	tttgtcttct	ataatatttc	ctttttttcc	cctctttgaa	tctctgtata	180
tttgattctt	aactaaaatt	gttctcttaa	atattctgaa	tcctggtaat	taaaagtttg	240
ggtgtatttt	ctttacctcc	aaggaaagaa	ctactagcta	caaaaaatat	tttggaataa	300
gcattgtttt	ggtataaggt	acatatattg	ggtgaagaca	ccagactgaa	gtaaacagct	360
gtgcatccaa	ttttattatg	ttttgtaagt	aacaatatgt	aatcaaaactt	ctaggtgact	420
tgagagtggg	acctcctata	tcattattta	gcaccgtttg	tgacagtaa		469

<210> 105

<211> 744

<212> DNA

<213> Human

<400> 105

ggcctgggac	aggattgagg	tatgttgacg	cctccagggc	ctgggggtctc	ctgcatgaag	60
aataccccctc	cccatttgac	tgtgaacttt	ttggcctgga	ttctggagaa	cagatttcca	120
ggattgtcag	ccagaaggca	gacagatgca	ggcacctacc	aagacctgac	ctcaggaagt	180
ggccctgccc	tacagcccag	ttgtcagcc	agggctgaag	gccatggggc	cccagcacc	240
ttgcttcagt	gccagcccct	ggaaggaacc	tcacaacagg	gatacagcaa	ggacactcca	300
gttccccag	tcctgccatg	gtgctaccct	gagggacagg	gatggagaca	gggcagccag	360
gtttgccagg	acctgcatag	cgggcccagg	actgcccttc	ctcttaagtc	atgccaaagc	420
ctccctgccc	agtctgagac	agtcgctggc	agggtgaccac	gacctgcgtg	gccctcccgg	480
cagttgtcat	ggtggttgta	ccccacccca	tccccctgag	gagacatggg	ctcagtccca	540
tgcttgggtgc	ccacagccac	aaagatggcc	atgggtctct	agcctgatat	tcgtggcctg	600
gcaggggtca	gcacccctga	gggcatccaa	gccatggtca	gaggaaagtg	ttggcaggct	660

cggcacagcc aaagaagtca ggacccacga gacgggggaa gccttccaga gccttcacct 720
tcacagggtc aaacttccag taga 744

<210> 106
<211> 401
<212> DNA
<213> Human

<400> 106
acattgttag gtgctgacct agacagagat gaactgaggt ccttgttttg tttgttcat 60
aatacaaagg tgctaattaa tagtatttca gatacttgaa gaatgttgat ggtgctagaa 120
gaatttgaga agaaatactc ctgtattgag ttgtatcgtg tgggtgattt tttaaaaaat 180
ttgatttagc attcatattt tccatcttat tcccaattaa aagtatgcag attatttggc 240
caaattctct tcagattcag catttgttct ttgccagtct cattttcatc ttcttccatg 300
gttccacaga agctttgttt cttgggcaag cagaaaaatt aaattgtacc tattttgtat 360
atgtgagatg tttaataaaa ttgtgaaaaa aatgaaataa a 401

<210> 107
<211> 1009
<212> DNA
<213> Human

<400> 107
cgagctatta tggtagcgaa ctttttttaa tgaggaattt catgatgatt taggaatttt 60
ctctcttggg aaaggcttcc cctgtgatga aaatgatgtg ccagctaaaa ttgtgtgcca 120
tttaaaaaact gaaaatattt taaaattatt tgtctatatt cttaaattgag ctttggatca 180
aactttaggc caggaccagc tcatgcgttc tcattcttcc ttttctcact ctttctctca 240
tcactcacct ctgtattcat tctgttgttt gggatagaaa aatcataaag agccaaccca 300
tctcagaacg ttgtggattg agagagacac tacatgactc caagtatatg agaaaaggac 360
agagctctaa ttgataactc tgtagttcaa aaggaaaaga gtatgcccaa ttctctctac 420
atgacatatt gagatttttt ttaatcaact tttaagatag tgatgttctg ttctaaactg 480
ttctgtttta gtgaaggtag atttttataa aacaagcatg gggattcttt tctaaggtaa 540
tattaatgag aagggaaaaa agtatcttta acagctcttt gttgaagcct gtggtagcmc 600
attatgttta taattgcaca tgtgcacata atctattatg atccaatgca aatacagctc 660
caaaaatatt aaatgtatat atattttaaa atgcctgagg aaatacattt ttcttaataa 720
actgaagagt ctcagtatgg ctattaaaaa aattattagc ctcctgttgt gtggctgcaa 780
aacatcacia agtgaccggt cttgagacct gtgaactgct gccctgttta gtaaaaaaaa 840
ttaatgcatt tctagagggg gaatatctgc catccagtgg tggaaatgtg gagtaagaaa 900
gctgggtggtc tgcttctgtg ctgtatgcca gccttttgcc ttaagttgag aggaggtcaa 960
ctttagctac tgtcttttgt ttgagagcca tggcaaaaaa aaaaaaaaaa 1009

This Page Blank (uspio)

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.

This Page Blank (uspto)